









Periodontitis and cardiovascular diseases: Consensus report

Mariano Sanz¹  | Alvaro Marco del Castillo² | Søren Jepsen³  | Jose R. Gonzalez-Juanatey⁴ | Francesco D'Aiuto⁵ | Philippe Bouchard⁶ | Iain Chapple⁷ | Thomas Dietrich⁷ | Israel Gotsman⁸ | Filippo Graziani⁹  | David Herrera¹  | Bruno Loos¹⁰  | Phoebus Madianos¹¹  | Jean-Baptiste Michel¹² | Pablo Perel^{13,14} | Burkert Pieske^{15,16} | Lior Shapira¹⁷  | Michael Shechter¹⁸ | Maurizio Tonetti¹⁹  | Charalambos Vlachopoulos²⁰ | Gernot Wimmer²¹

¹Department of Dental Clinical Specialties, ETEP Research Group, Faculty of Odontology, University Complutense of Madrid, Madrid, Spain

²Cardiology Department, Hospital Universitario Ramon y Cajal, Madrid, Spain

³Department of Periodontology, Operative and Preventive Dentistry, University of Bonn, Bonn, Germany

⁴Cardiology Department, University Hospital, IDIS, CIBERCV, University of Santiago de Compostela, Santiago de Compostela, Spain

⁵Department of Periodontology, Eastman Dental Institute and Hospital, University College London, London, UK

⁶U.F.R. d'odontologie, Université Paris Diderot, Hôpital Rothschild AP-HP, Paris, France

⁷School of Dentistry, Institute of Clinical Sciences, College of Medical & Dental Sciences, The University of Birmingham, Birmingham, UK

⁸Heart Institute, Hadassah University Hospital, Jerusalem, Israel

⁹Department of Surgical, Medical and Molecular Pathology and Critical Care Medicine, University of Pisa, Pisa, Italy

¹⁰ACTA University, Amsterdam, The Netherlands

¹¹Department of Periodontology, School of Dentistry, National and Kapodistrian University of Athens, Athens, Greece

¹²Inserm Unit 1148, Laboratory for Translational CV Science, X. Bichat Hospital, Paris, France

¹³World Heart Federation, Geneva, Switzerland

¹⁴Centre for Global Chronic Conditions, London School of Hygiene & Tropical Medicine, London, UK

¹⁵Department of Internal Medicine & Cardiology, Charité Universitätsmedizin Berlin, Berlin, Germany

¹⁶DZHK (German Center for Cardiovascular Research) Partnersite Berlin, German Heart Institut Berlin, Berlin, Germany

¹⁷Department of Periodontology, Hebrew University – Hadassah Faculty of Dental Medicine, Jerusalem, Israel

¹⁸Leviev Heart Center, Chaim Sheba Medical Center, Tel Hashomer and the Sackler Faculty of Medicine, Tel Aviv University, Tel Aviv-Yafo, Israel

¹⁹Department of Periodontology, Prince Philip Dental Hospital, The University of Hong Kong, Hong Kong, Hong Kong

²⁰1st Department of Cardiology, National and Kapodistrian University of Athens, Athens, Greece

²¹Department of Prosthetic Dentistry, School of Dental Medicine, Karl-Franzens University Graz, Graz, Austria

Correspondence

Mariano Sanz, Department of Dental Clinical Specialties and ETEP Research Group, Faculty of Odontology, University Complutense of Madrid, Plaza Ramon y Cajal, E-28040 Madrid, Spain.
Email: marianosanz@odon.ucm.es

Funding information

DENTAID SA

Abstract

Background: In Europe cardiovascular disease (CVD) is responsible for 3.9 million deaths (45% of deaths), being ischaemic heart disease, stroke, hypertension (leading to heart failure) the major cause of these CVD related deaths. Periodontitis is also a chronic non-communicable disease (NCD) with a high prevalence, being severe periodontitis, affecting 11.2% of the world's population, the sixth most common human disease.

Material and Methods: There is now a significant body of evidence to support independent associations between severe periodontitis and several NCDs, in particular CVD. In 2012 a joint workshop was held between the European Federation of

Periodontology (EFP) and the American Academy of Periodontology to review the literature relating periodontitis and systemic diseases, including CVD. In the last five years important new scientific information has emerged providing important emerging evidence to support these associations

Results and Conclusions: The present review reports the proceedings of the workshop jointly organised by the EFP and the World Heart Federation (WHF), which has updated the existing epidemiological evidence for significant associations between periodontitis and CVD, the mechanistic links and the impact of periodontal therapy on cardiovascular and surrogate outcomes. This review has also focused on the potential risk and complications of periodontal therapy in patients on anti thrombotic therapy and has made recommendations for dentists, physicians and for patients visiting both the dental and medical practices.

KEYWORDS

anti thrombotic therapy, atherosclerosis, bacteremia, cardiovascular disease, chronic inflammation, periodontal therapy, periodontitis

1 | INTRODUCTION

Non-communicable diseases (NCDs) are rising in prevalence globally in line with an increasingly ageing population, refined diets and sedentary lifestyles and account for 41 million deaths each year, or 71% of all global deaths (G. B. D. Risk Factors Collaborators, 2016). Approximately 80% of people over 65-years of age in the United States are affected by one or more NCDs and 77% exhibit at least two NCDs, creating a significant burden of disease to individuals and to the healthcare economy (Centres for Disease Control & Prevention, 2011). The comorbid presence of two or more NCDs presents a major challenge to the economy, equating to two-thirds of all health costs in the United States (Centres for Disease Control & Prevention, 2013); however, <1% USA health expenditure is focussed on prevention to improve overall health (U.S. Senate Committee on Health, 2011).

The greatest global NCD burden arises due to cardiovascular disease (CVD), responsible for 17.9 million deaths (a third of total mortality), and 45% of NCD-induced mortality (Roth et al., 2017). In Europe, CVD is responsible for 3.9 million deaths (45% of deaths), and whilst CVD mortality rates are reducing, the absolute numbers have increased in the last 25 years, due to an increasingly ageing population (Wilkins et al., 2017). Ischaemic heart disease, stroke, hypertension (leading to heart failure), rheumatic heart disease, cardiomyopathy and atrial fibrillation cause over 95% of CVD-related deaths (Roth et al., 2015).

In this consensus report, the term CVD is used as a general term for atherosclerotic diseases, principally coronary heart disease, cerebrovascular disease and peripheral vascular disease. A number of chronic infectious, inflammatory and immune diseases are associated with significantly higher risks of adverse cardiovascular events, including rheumatoid arthritis, psoriasis, systemic lupus erythematosus and periodontitis (Roth et al., 2015), consistent with the thesis that chronic elevations in the systemic inflammatory burden are causally related to CVD development and its sequelae. Whilst there

is evidence for over 50 gene polymorphisms playing a role in the modulation of atherogenesis (Holdt & Teupser, 2015), effect sizes are small and the major traditional risk factors for CVD remain the lifestyle factors, principally tobacco smoking, dyslipidaemia, hypertension and altered glucose metabolism. The latter correlate strongly with diets high in saturated fats, salt and refined sugars and contribute to obesity and type 2 diabetes mellitus, major attributable risk factors for myocardial infarction (Joseph et al., 2017). The same risk factors account for over 90% of the stroke burden (O'Donnell et al., 2016), yet all are modifiable through improved lifestyles including reducing salt, saturated fat and refined carbohydrate intake, exercising, increasing intake of antioxidant micronutrients and regular moderate alcohol consumption (Joseph et al., 2017).

Periodontitis is also a NCD with a high prevalence of 45%–50% overall, with the most severe form affecting 11.2% of the world's population, being the sixth most common human disease (Kassebaum et al., 2014). The Global Burden of Diseases, Injuries, and Risk Factors Study (2017) of years lost to disability (YLD) reported that from 1990 to 2017 oral diseases (mainly periodontitis and caries) contributed the most YLD in age-standardized prevalence rates from 354 diseases and injuries across 195 countries (G. B. D. Disease Injury & Incidence & Prevalence Collaborators, 2018). There is now a significant body of evidence to support independent associations between severe periodontitis and several NCDs including diabetes (Chapple, Genco, & Working group 2013 of the joint EFP/AAP Workshop, 2013), cardiovascular disease (Tonetti et al., 2013), chronic obstructive pulmonary disease (Linden, Lyons, & Scannapieco, 2013) and chronic kidney disease (CKD) (Sharma, Dietrich, Ferro, Cockwell, & Chapple, 2016). Indeed, severe periodontitis is independently and significantly associated with all-cause and cardiovascular mortality in several different populations (Linden et al., 2012; Sharma et al., 2016). Proposed mechanisms include bacteraemia and the associated systemic inflammatory sequelae, including elevations in C-reactive protein and oxidative

stress (Schenkein & Loos, 2013). In populations with multimorbidity, for example chronic kidney disease with comorbid diabetes and periodontitis, periodontitis is associated with significantly reduced survival from all-cause and cardiovascular mortality (Sharma et al., 2016). It appears therefore that periodontitis may be a modifiable non-traditional risk factor for CVD.

In 2012, a joint workshop was held between the European Federation of Periodontology (EFP) and the American Academy of Periodontology to review the literature relating periodontitis and systemic diseases, including CVD. The consensus report was based upon four technical papers that systematically reviewed the evidence for epidemiological associations between periodontitis and incident CVD (Dietrich, Sharma, Walter, Weston, & Beck, 2013), mechanisms of biological plausibility relating to periodontal bacteria and systemic inflammation (Reyes, Herrera, Kozarov, Roldan, & Progulski-Fox, 2013; Schenkein & Loos, 2013) and periodontal intervention studies (D'Aiuto, Orlandi, & Gunsolley, 2013). The workshop concluded that there was consistent and strong epidemiological evidence that periodontitis imparts increased risk for future atherosclerotic cardiovascular disease. It also concluded that the impact of periodontitis on CVD was biologically plausible, via translocated circulating oral microbiota, which may directly or indirectly induce systemic inflammation that impacts upon the development of atherothrombogenesis, and whilst *in vitro*, pre-clinical and clinical studies supported the interaction and associated biological mechanisms, intervention trials were not sufficiently adequate to draw further conclusions at that time.

The present workshop was jointly organized by the EFP and the World Heart Federation (WHF) to include global experts in both periodontal and cardiovascular disciplines and was held in Madrid on 18th and 19th February 2019. Four technical reviews updating the evidence base from the 2012 workshop were prepared and supplemented by additional studies discussed at the workshop. The reviews focussed on epidemiological associations (Herrera, Molina, Buhlin, & Klinge, 2019), mechanistic links (Schenkein, Papapanou, Genco, & Sanz, 2019), results from intervention studies (Orlandi, Graziani, & D'Aiuto, 2019) and the potential risk and complications of periodontal therapy in patients undertaking antithrombotic (antiplatelet and anticoagulant) therapy.

Whilst this consensus report focuses predominantly on relevant evidence published since the 2012 workshop, there are biological areas that have subsequently come to prominence, where the underpinning body of evidence was not covered in the 2013 consensus report, and hence, certain pre-2012 manuscripts are referenced to ensure the context of these recent studies is clear.

Furthermore, section 4.3 "*What is the effect of statin intake on clinical periodontal outcomes?*" and section 5 "*Cardiovascular risks and complications of periodontal therapeutic interventions*" were not dealt in the previous workshop, and hence, a full appraisal of the scientific evidence was carried out in this consensus meeting.

Finally, following the review of the presented evidence, recommendations for both medical and dental teams, as well as patients and the public, were elaborated.

2 | EPIDEMIOLOGIC EVIDENCE ON THE ASSOCIATION BETWEEN PERIODONTITIS AND CVD

2.1 | Do people with periodontitis have a higher prevalence of subclinical cardiovascular disease?

There is evidence from epidemiological studies that periodontitis patients exhibit significant endothelial dysfunction, measured by flow-mediated dilation (FMD), arterial stiffness (e.g. pulse wave velocity—PWV) and a significantly greater thickness of the carotid intima-media (cIMT) and elevated arterial calcification scores. There is one imaging study (ATHEROREMO-IVUS study) associating high levels of antibodies against periodontal pathogens and a lower extent of positive atheromatous plaque remodelling (de Boer et al., 2014).

2.2 | Do people with periodontitis have a higher prevalence of coronary artery disease and risk of myocardial infarction and other coronary events?

There is robust evidence from epidemiological studies for a positive association between periodontitis and coronary heart disease. A systematic review (Dietrich et al., 2013), which was updated in preparation for this workshop, identified a total of 6 case-control and cohort studies epidemiological studies, published in the last five years, which demonstrated an increased risk of a first coronary event in patients with clinically diagnosed periodontitis or more severe periodontitis compared to patients without periodontitis or less severe periodontitis. Relative risk estimates vary between studies, depending on population characteristics and periodontitis case definitions. There are two cohort studies reporting an association between periodontitis and higher cardiovascular mortality (due to coronary heart disease and cerebrovascular disease).

2.3 | Do people with periodontitis have a higher prevalence of cerebrovascular disease and risk of stroke?

There is evidence from epidemiologic studies for a positive association between periodontitis and cerebrovascular disease. A systematic review (Dietrich et al., 2013), which was updated in preparation for this workshop, identified a total of three case-control and cohort studies, which demonstrate an increased risk of a first cerebrovascular event in patients with clinically diagnosed periodontitis or more severe periodontitis compared to patients without periodontitis or less severe periodontitis. Relative risk estimates vary between studies, depending on population characteristics and periodontitis case definitions. Furthermore, a recent analysis of data from the ARIC study demonstrated an association between periodontal profile class and incident ischaemic stroke. In this cohort, patients with periodontitis had more than double the risk of cardioembolic and thrombotic stroke compared with periodontally healthy individuals (Sen et al., 2018). In addition, as previously documented, there are two cohort studies reporting an association between periodontitis

and higher cardiovascular mortality (due to coronary heart disease and cerebrovascular disease) (Dietrich et al., 2013).

2.4 | Do people with periodontitis have a higher prevalence and incidence of Peripheral Artery Disease (PAD)?

There is limited but consistent evidence that individuals with periodontitis have a higher prevalence and incidence of PAD compared to individuals without periodontitis (Yang et al., 2018). For cross-sectional data, the most significant evidence comes from two large, population-based studies in the United States (NHANES 1999–2002) and South Korea (KoGES-CAVAS). Both studies found a positive association between the extent of clinical attachment loss (NHANES 1999–2002) and severity of radiographic bone loss (KoGES-CAVAS) with PAD, defined using the Ankle Brachial Index (ABI), with adjusted odds ratios (OR) of 2.2 (95% confidence interval [1.2; 2.4]) and 2.0 (95% CI [1.1; 3.9]), respectively (Ahn et al., 2016; Lu, Parker, & Eaton, 2008). One prospective cohort study conducted in male veterans in the United States reported a positive association between periodontitis (measured by severity of radiographic bone loss) and the incidence of PAD over a 25- to 30-year follow-up period, with an adjusted OR of 2.3 (95% CI [1.3; 3.9]) (Mendez et al., 1998). There are no studies that have evaluated the association between periodontitis and the incidence of Major Adverse Limb Events (MALE).

2.5 | Do people with periodontitis have a higher risk of other CVDs or conditions (heart failure, atrial fibrillation)?

Several studies report positive associations between periodontitis and heart failure. There is evidence from a large Asian study using the Taiwanese National Health Insurance Research Database reporting a significantly higher incidence of atrial fibrillation in individuals with periodontal diseases compared to individuals without periodontal diseases (hazard ratio—HR = 1.31, 95% CI [1.25, 1.36]) (Chen, Lin, Chen, & Chen, 2016).

2.6 | Do people with a history of cardiovascular disease have a different incidence or progression of periodontitis?

There is currently limited scientific evidence that CVD is a risk factor for the onset or progression of periodontitis.

2.7 | Do people with periodontitis with history of cardiovascular disease have a higher chance of experiencing a subsequent event?

From three studies investigating the association between periodontitis and secondary cardiovascular events, two large studies did not find a significant association (Dorn et al., 2010; Reichert et al., 2016); however, a small study (100 subjects) reported a significant association (HR = 2.8, 95% CI [1.2; 6.5]) with recurrent cerebrovascular events (Sen et al., 2013).

3 | MECHANISMS THAT MAY EXPLAIN THE EPIDEMIOLOGICAL ASSOCIATIONS BETWEEN PERIODONTITIS AND CVD

3.1 | Is there evidence of a higher incidence of bacteremia following oral function/intervention in periodontitis patients compared to periodontally healthy subjects?

There is evidence that oral bacterial species can enter the circulation and cause bacteremia, which has been demonstrated following daily life activities (toothbrushing, flossing, chewing or biting an apple), although it has been studied more frequently following professional interventions (tooth polishing, scaling, tooth extraction, surgical extraction of third molars and periodontal probing).

The risk of bacteremia has been associated with periodontal health status in a systematic review, suggesting a higher risk of bacteremia associated with gingival inflammation (Tomas, Diz, Tobias, Scully, & Donos, 2012). A recent randomized clinical trial (RCT) concluded that periodontal therapy (by means of scaling and root planing, SRP) induced bacteremia in both gingivitis and periodontitis patients, but the magnitude and frequency were greater among periodontitis patients (Balejo et al., 2017).

Whilst there are methodological limitations in some of the reported studies, the overall picture supports the contention that bacteremia results from daily life activities and oral interventions, and it is more frequent of longer duration and involves more virulent bacteria in periodontitis patients.

3.2 | Is there evidence for the presence of oral bacteria in atheroma lesions?

There is evidence through traces of DNA, RNA or antigens derived from oral bacterial species, mainly periodontal pathogens, that have been identified in atherothrombotic tissues. Studies have attempted to correlate the presence of these bacteria in atherothrombotic tissues, with other sample sources (subgingival plaque, serum, etc.), in the same patients, and these suggest that in periodontitis patients there is a higher probability of a positive correlation (Armingohar, Jørgensen, Kristoffersen, Abesha-Belay, & Olsen, 2014; Mahendra, Mahendra, Felix, & Romanos, 2013). At least two studies have demonstrated viable *P. gingivalis* and *A. actinomycetemcomitans* in atherothrombotic tissue when culturing the atheroma samples (Kozarov, Dorn, Shelburne, Dunn, & Progulske-Fox, 2005; Rafferty et al., 2011).

3.3 | Do we have evidence that periodontal bacteria and/or bacterial products and virulence factors influence the pathophysiology of atherosclerosis?

Different animal models have been employed to provide evidence that periodontal pathogens can promote atheroma formation. *P. gingivalis* has been shown to accelerate atherosclerosis in murine

models, to induce fatty streaks in the aorta of rabbits and to induce aortic and coronary lesions after bacteremia in normocholesterolaemic pigs (Schenkein & Loos, 2013).

Recently, further evidence has emerged using hyperlipidemic ApoE null mice after infection with *P. gingivalis* and also with a polymicrobial experimental infection (*P. gingivalis*, *Treponema denticola*, *Tannerella forsythia* and *Fusobacterium nucleatum*). A polymicrobial infection was shown to induce aortic toll-like receptor (TLR) and inflammasome signalling, with an enhanced oxidative stress reaction generated within the aortic endothelial cells (Chukkapalli et al., 2015; Velsko et al., 2014, 2015).

There is also *in vitro* evidence of intracellular entry by periodontal pathogens (*P. gingivalis*, *A. actinomycetemcomitans*, etc.) (Reyes et al., 2013). *In vivo* and *in vitro* studies demonstrate the importance of the fimbriae of *P. gingivalis* to host cell entry and to promote atherothrombotic lesions in experimental models (Yang et al., 2014). *In vitro* experiments have shown that certain bacterial strains expressing *P. gingivalis* hemagglutinin A (HagA) have an increased capability to adhere and enter human coronary artery endothelial cells (HCAEC) (Belanger, Kozarov, Song, Whitlock, & Progske-Fox, 2012).

3.4 | Do we have evidence that periodontitis patients exhibit increased production and/or levels of inflammatory mediators that also associated with the pathophysiology of atherosclerosis?

There is evidence of significantly higher levels of C-reactive protein (CRP) in periodontitis patients versus healthy controls and in CVD and periodontitis patients compared with either condition alone. The effect of periodontal therapy has been shown to associate with a significant decrease in CRP levels, along with improvements in surrogate measurements of cardiovascular health (Demmer et al., 2013; Koppolu et al., 2013; Patil & Desai, 2013).

There is evidence of elevated levels of serum interleukin (IL)-6 in periodontitis patients and lower levels of IL-4 and IL-18. The effect of periodontal therapy has shown a significant decrease in the serum levels of IL-6, serum amyloid A and alpha 1 anti-chymotrypsin. Peripheral neutrophils from periodontitis patients release excess IL-1 β , IL-8, IL-6 and tumour necrosis factor (TNF)- α when stimulated by periodontal pathogens. Periodontal therapy only partially reduces the cytokine hyper-reactivity with some evidence of a constitutively elevated response (Ling, Chapple, & Matthews, 2016).

3.5 | Do we have evidence that periodontitis patients develop elevations in thrombotic factors that are also associated with the pathophysiology of atherothrombosis?

There is evidence of significantly higher levels of fibrinogen in periodontitis patients versus healthy controls, and in CVD and periodontitis patients compared with either condition alone (Chandy et al., 2017). Periodontal therapy appears to result in a significant decrease

in fibrinogen levels (Lopez et al., 2012; Vidal, Cordovil, Figueredo, & Fischer, 2013).

There is evidence from different studies of significantly higher levels of platelet activation markers in periodontitis patients and that these higher levels may be reversed by periodontal therapy (Arvanitidis, Bizzarro, Alvarez Rodriguez, Loos, & Nicu, 2017). However, there is conflicting evidence that significantly higher levels of plasminogen activator inhibitor (PAI) are found in periodontitis patients (Schenkein & Loos., 2003).

3.6 | Do we have evidence that periodontitis patients demonstrate elevated serum antibody levels that cross-react with antigens in cardiovascular tissues?

There is evidence that HSPs from periodontal pathogens (*Porphyromonas gingivalis*, *Tannerella forsythia*, *Aggregatibacter actinomycetemcomitans* and *Fusobacterium nucleatum*) generate antibodies that can cross-react with human HSPs. These antibodies have been shown to activate cytokine production, as well as monocyte and endothelial cell activation.

The presence of anti-cardiolipin antibodies has been significantly associated with periodontitis patients, which reversed following periodontal therapy. There is some evidence that periodontal pathogens can elicit antibodies that cross-react with cardiolipin (Schenkein & Loos., 2003).

In three out of four population-based studies (Parogene study, NHANES III, DANHES), higher levels of serum immunoglobulin (Ig)G against *P. gingivalis* were associated with periodontitis patients and cardiovascular disease (acute coronary syndrome, death from cardiovascular disease and cardiovascular disease). The ATHEROREMO-IVUS study failed to demonstrate an association between serum levels of IgG and IgA against *P. gingivalis*, *A. actinomycetemcomitans*, *T. forsythia* and *P. intermedia* and major adverse cardiac events (MACE) (de Boer et al., 2014). This is consistent with data from Boillot et al. (2016).

3.7 | Do we have evidence that periodontitis patients exhibit dyslipidaemia?

There is evidence from systematic reviews that serum total cholesterol levels, low-density lipoproteins (LDL), triglycerides, very-low-density lipoproteins (VLDL), oxidized LDL and phospholipase A2 are elevated in periodontitis. High-density lipoprotein (HDL) levels are reduced in periodontitis patients compared with controls (Schenkein & Loos., 2003). These levels are reversed after periodontal therapy (Teeuw et al., 2014).

3.8 | Do we have evidence for peripheral blood neutrophil hyper-responsiveness in reactive oxygen species and protease production in periodontitis patients?

There is strong mechanistic evidence that peripheral blood neutrophils (PBNs) from periodontitis patients produce higher levels of total

and extracellular reactive oxygen species (ROS) than healthy controls, under various conditions of priming and stimulation and from unstimulated cells (Ling et al., 2016; Matthews, Wright, Roberts, Cooper, & Chapple, 2007a). This hyper-reactivity to stimulation by periodontal bacteria is reduced following successful periodontal therapy to control patient levels, but the unstimulated hyperactivity remains, suggesting that constitutive and reactive mechanisms underlie neutrophil hyper-responsiveness in periodontitis (Matthews, Wright, Roberts, Ling-Mountford, et al., 2007b). Gene expression data in PBNs support the functional data (Wright, Matthews, Chapple, Ling-Mountford, & Cooper, 2008). Serum antioxidant levels and those in gingival crevicular fluid (GCF) are reduced in periodontitis patients, reflecting increased ROS activity (Chapple, Brock, Milward, Ling, & Matthews, 2007). These data are supported by a study of endarterectomy samples, which demonstrated evidence for activation of the ROS-generating systems in neutrophils, specifically the presence of myeloperoxidase (MPO), cell-free DNA and DNA-MPO complexes (Range et al., 2014).

3.9 | Are there common genetic risk factors between periodontitis and CVDs?

There is scientific evidence of pleiotropy between periodontitis and cardiovascular diseases (Aarabi et al., 2017; Munz et al., 2018; Schaefer et al., 2015, 2011). The highly pleiotropic genetic locus *CDKN2B-AS1* (chromosome 9, p21.3) associated with coronary artery disease, type 2 diabetes, ischaemic stroke and Alzheimer's disease is also consistently associated with periodontitis (Aarabi et al., 2017; Ernst et al., 2010; Loos, Papantonopoulos, Jepsen, & Laine, 2015; Munz et al., 2018). Its function appears to be related to the regulation of gene expression (Hubberten et al., 2019). Interestingly, a pilot study identified that a genetic variant in the *CDKN2B-AS1* locus was associated with the extent of elevated levels of C-reactive protein in periodontitis (Teeuw, Laine, Bizzarro, & Loos, 2015).

A conserved non-coding element within *CAMTA1* upstream of *VAMP3*, also first identified as a genetic susceptibility locus for coronary artery disease, was found to be associated with periodontitis (Schaefer et al., 2015). A GWAS suggested that the *VAMP3* locus was associated with a higher probability of subgingival overgrowth of periodontal pathogens (Divaris et al., 2012).

There is evidence for plasminogen (PLG) as a shared genetic risk factor for coronary artery disease and periodontitis (Schaefer et al., 2015).

The 4th pleiotropic locus between coronary artery disease and periodontitis is a haplotype block at the *VAMP8* locus (Munz et al., 2018).

These shared genetic factors suggest a mechanistic link or immunological commonalities between coronary artery disease and periodontitis. The impairment of the regulatory pathways by genetic factors may be a common pathogenic denominator of at least coronary artery disease and periodontitis. There are indications that aberrant inflammatory reactivity, determined by genetic variants in the loci *CDKN2B-AS1* (*ANRIL*), *PLG*, *CAMTA1/VAMP3* and *VAMP8* could

partially explain the epidemiological link between periodontitis and cardiovascular diseases.

4 | EVIDENCE FROM INTERVENTION STUDIES

4.1 | Is there an effect of periodontitis treatment in preventing or delaying ACVD events?

4.1.1 | Primary prevention

There have been no prospective randomized controlled periodontal intervention studies on primary prevention of cardiovascular diseases (including first ischaemic events or cardiovascular death) since the last consensus report (Tonetti et al., 2013). The Group questioned the feasibility of performing adequately powered RCTs in primary prevention at a population level due to important ethical, methodological and financial considerations.

However, consistent observational evidence suggests that several oral health interventions including self-performed oral hygiene habits (toothbrushing) (two studies (de Oliveira, Watt, & Hamer, 2010; Park et al., 2019)), dental prophylaxis (one study (Lee, Hu, Chou, & Chu, 2015)), increased self-reported dental visits (one study (Sen et al., 2018)) and periodontal treatment (three studies (Holmlund, Lampa, & Lind, 2017; Lee et al., 2015; Park et al., 2019)) produced a reduction in the incidence of ACVD events.

Cross-sectional data of The Scottish Health Surveys from 1995 to 2003 pertaining 11,869 men and women (mean age of 50 years) were linked to a database of hospital admissions and deaths with follow-up until December 2007 (Information Services Division, Edinburgh) (de Oliveira et al., 2010). Participants who brushed less than once a day exhibited the highest incidence of ACVD events (HR = 1.7, 95% CI [1.3; 2.3]) compared with those who brushed twice a day, indicating that self-performed oral hygiene routines may reduce the incidence of ACVD.

A retrospective nationwide, population-based study in Taiwan, including 511,630 participants with periodontitis and 208,713 controls, used the Longitudinal Health Insurance Database 2000 to estimate the incidence rate of ACVD events from 2000 to 2015 (Lee et al., 2015). The hazard ratio for acute myocardial infarction was reduced more in the group of periodontitis patients who received dental prophylaxis (HR = 0.90, 95% CI [0.86; 0.95]) than intensive treatment (including gingival curettage, scaling and root planing, and/or periodontal flap operation and/or tooth extraction) (HR = 1.09, 95% CI [1.03; 1.15]). Consistent reductions in the incidence rate of ischaemic stroke were observed in both the dental prophylaxis (HR = 0.78, 95% CI [0.75; 0.91]) and intensive treatment groups (HR = 0.95, 95% CI [0.91; 0.99]).

A cohort of 8,999 patients with periodontitis who received a complete (non-surgical and if needed surgical) periodontal treatment protocol was followed between 1979 and 2012 (Holmlund et al., 2017). During the study follow-up, poor responders to the periodontal treatment had an increased incidence of ACVD events (incidence rate -IR = 1.28, 95% CI [1.07; 1.53]) compared with good

responders, suggesting that successful periodontal treatment could reduce the incidence of ACVD events.

In the Atherosclerosis Risk in Communities (ARIC) study including 6,736 participants followed during 15 years, self-reported regular dental care users had a lower risk for ischaemic stroke (HR = 0.77, 95% CI [0.63; 0.94]) compared with episodic care users (Sen et al., 2018).

In a prospective population-based study using data from the National Health Insurance System-National Health Screening Cohort (NHISHEALS) including 247,696 participants free from any CVD history recruited between 2002 and 2003, reported that an increased number of dental caries lesions, the presence of periodontitis and a greater loss of teeth were all associated with an increased risk of future major cardiovascular events (MACEs), including cardiovascular death, acute myocardial infarction, heart failure, and stroke (Park et al., 2019). One additional toothbrushing episode per day was associated with a reduced incidence of ACVD events (HR = 0.91, 95% CI [0.89, 0.93]) and regular professional cleaning reduced the risk even further (HR = 0.86, 95% CI [0.82; 0.90]).

In summary, progression of ACVD may be influenced by successful periodontal treatment independent of traditional CVD risk factor management.

4.1.2 | Secondary prevention

There is only one pilot multicentre study on secondary prevention of ACVD events (PAVE (Couper et al., 2008; Offenbacher et al., 2009)), which reported no statistically significant difference in the rate of CVD events between patients who underwent treatment of periodontitis versus community care (risk ratio -RR = 0.72, 95% CI [0.23; 2.22]). Several methodological limitations highlighted in the trial limit the applicability/usefulness of such evidence to inform the research and healthcare communities.

Thus, there is insufficient evidence to support or refute the potential benefit of the treatment of periodontitis in preventing or delaying ACVD events (Li et al., 2017).

4.2 | What is the effect of the treatment of periodontitis in improving surrogate parameters of CVD?

Table 1 summarizes the evidence on the effect of periodontal therapy on surrogate markers of CVD. There is moderate evidence for reduction of low-grade inflammation as assessed by serum levels of CRP, IL-6 and improvements in surrogate measures of endothelial function (flow-mediated dilatation of the brachial artery).

Moderate evidence suggests that periodontal treatment does not have an effect on lipid fractions whilst there is limited evidence, suggesting that periodontal treatment reduces arterial blood pressure and stiffness, subclinical ACVD (as assessed by mean carotid intima-media thickness) and insufficient evidence of an effect on ACVD biomarkers of coagulation, endothelial cell activation and oxidative stress.

4.3 | What is the effect of statin intake on clinical periodontal outcomes?

Statins are medications prescribed to decrease LDL cholesterol. Numerous trials have demonstrated their benefit for the prevention of cardiovascular diseases (Yebo, Aschmann, Kaufmann, & Puhon, 2019).

Interestingly, statins possess various additional properties relevant to the pathogenesis and treatment of periodontitis (Estanislau et al., 2015). In particular, it has been reported that statins are anti-inflammatory (Koh et al., 2002; Paumelle et al., 2006; Quist-Paulsen, 2010; Rosenson, Tangney, & Casey, 1999; Sakoda et al., 2006) can promote bone formation (Garrett, Gutierrez, & Mundy, 2001; Liu et al., 2012; Mundy et al., 1999; Viereck et al., 2005), can inhibit matrix metalloproteinases (MMPs) (Koh et al., 2002; Luan, Chase, & Newby, 2003; Poston et al., 2016) and possess anti-microbial properties (Ting, Whitaker, & Albandar, 2016).

A systematic review with meta-analysis of pre-clinical in vivo trials reported a positive effect of local or systemic statin administration for the prevention of alveolar bone loss in experimental periodontitis models in rodents (Bertl et al., 2018).

Several observational clinical studies have evaluated the effect of systemic statin intake on periodontal conditions (Lindy, Suomalainen, Mäkelä, & Lindy, 2008; Meisel, Kroemer, Nauck, Holtfreter, & Kocher, 2014; Sangwan, Tewari, Singh, Sharma, & Narula, 2013; Saver, Hujoel, Cunha-Cruz, & Maupome, 2007; Saxlin, Suominen-Taipale, Knuuttila, Alha, & Ylostalo, 2009; Subramanian et al., 2013). Statin use was not found to be associated with decreased tooth loss in adults with chronic periodontitis when analysing administrative health plan data (Saver et al., 2007). However, a 5-year population-based follow-up study comparing participants treated with statins with those who did not medicate with statins concluded that long-term treatment with statins was associated with reduced tooth loss (Meisel et al., 2014). Furthermore, patients on statin medication were reported to exhibit significantly fewer signs of periodontal inflammatory lesions than patients without a statin regimen (Lindy et al., 2008). A cross-sectional study compared the periodontal status of patients with hyperlipidaemia (with or without statin intake) to normolipidaemic individuals and found higher gingival bleeding and probing depths in the hyperlipidaemic patients who were not statin users (Sangwan et al., 2013). In a RCT, periodontal patients with risk factors or with established atherosclerosis were assigned to either high- or low-dose statin intake (Subramanian et al., 2013). After 3 months, a significant reduction of periodontal inflammation was seen in the high-dose compared to the low-dose group. Thus, within the limits of the above-reported studies, there is some limited evidence, suggesting that statins may have a positive impact on periodontal health.

Very few clinical studies have been designed to evaluate the effect of adjunctive systemic statin intake in conjunction with periodontal therapy (Fajardo, Rocha, Sanchez-Marin, & Espinosa-Chavez, 2010; Fentoglu et al., 2012; Sangwan, Tewari, Singh, Sharma, & Narula, 2016). In a randomized placebo-controlled pilot study in 38 patients with chronic periodontitis, adjunctive statin intake led to beneficial effects on radiological bone loss and tooth mobility after 3 months

TABLE 1 Summary of the evidence on the effect of periodontal therapy on surrogate markers of cardiovascular diseases

Topic	Outcome	Number of RCTs and SR since last consensus	References	Effect	Overall Level of Evidence
Effect of Periodontal Therapy on Lipids	Lipids (multiple)	6 RCTs	Caula, Lira-Junior, Tinoco, and Fischer (2014); D'Aiuto et al. (2018); Deepti, Tewari, Narula, Singhal, and Sharma (2017); Fu, Li, Xu, Gong, and Yang (2016); Hada, Garg, Ramteke, and Ratre (2015); Kapellas et al. (2014)	No	Moderate
Effect of Periodontal Therapy on Blood Pressure	Systolic, diastolic	3 RCTs	D'Aiuto et al. (2018); Hada et al. (2015); Zhou et al. (2017)	Yes	Limited
Effect of Periodontal Therapy on Endothelial Function	Endothelial Function (multiple measures)	2 RCTs 1 SR	D'Aiuto et al. (2018); Saffi et al. (2018) Steffel et al. (2018b)	Yes	Moderate
Effect of Periodontal Therapy on interleukin (IL)-6	IL-6	3 RCTs	Fu et al. (2016); Kapellas et al. (2014); Zhou et al. (2017)	Yes	Moderate
Effect of Periodontal Therapy on C-Reactive Protein (CRP)	CRP	5 SR 7 RCTs following 2014	Demmer et al. (2013); Freitas et al. (2012); Ioannidou, Malekzadeh, and Dongari-Bagtzoglou (2006); Paraskevas, Huizinga, and Loos (2008); Teeuw et al. (2014) D'Aiuto et al. (2018); Deepti et al. (2017); Kaushal, Singh, Lal, Das, and Mahdi (2019); Caula et al. (2014); Hada et al. (2015); Kapellas et al. (2014); Zhou et al. (2017)	Yes	Moderate
Effect of Periodontal Therapy on Pulse Wave Velocity (PWV)	PWV	1 RCT	Kapellas et al. (2014)	No	Limited
Effect of Periodontal Therapy on carotid intima-media thickness (cIMT)	Common cIMT	1 RCT	Kapellas et al. (2014)	Yes	Limited

Abbreviation: RCT, randomized clinical trial; SR, systematic review.

(Fajardo et al., 2010). Another 3-month study compared the treatment response to nonsurgical periodontal therapy in 107 chronic periodontitis patients (35 normolipidaemic as control, 36 hyperlipidaemic on non-pharmacological therapy and 36 hyperlipidaemic on statins) and found a greater improvement in gingival index in the normolipidaemic control and in the statin groups (Sangwan et al., 2016). Based on this limited evidence, two recent systematic reviews with meta-analysis on the effects of (local and systemic) statins on periodontal treatment concluded that systemic statin intake does not enhance the outcomes of periodontal therapy (Bertl et al., 2017; Muniz et al., 2018).

5 | CARDIOVASCULAR RISKS AND COMPLICATIONS OF PERIODONTAL THERAPEUTIC INTERVENTIONS

5.1 | Is there an ischaemic cardiovascular risk for patients undergoing periodontal therapy?

Non-surgical treatment of periodontitis involving supra- and subgingival instrumentation of the affected dentition (under local anaesthesia) is often delivered in several short sessions. Alternatively,

full-mouth non-surgical periodontal treatment can be performed within 24 hours.

Delivering periodontal treatment in a full-mouth fashion (i.e. within 24 hours) triggers a one-week acute systemic inflammatory response associated with transient impairment of endothelial function (Orlandi et al., 2019). This distant effect is not observed when periodontal treatment is delivered across several separate sessions (Graziani et al., 2015). This is achieved by limiting the number of teeth involved and the time devoted to completing the dental instrumentation. These findings raise the question of whether performing longer sessions of periodontal treatment could contribute to an individuals' inflammatory burden/risk and increase their short-term risk of suffering from a vascular event. There is consistent and strong observational evidence that common acute infections/inflammatory responses are associated at a population level with an increased risk of vascular events within the first 4 weeks of the infectious/inflammatory event (Smeeth et al., 2004).

5.1.1 | At population level

There is no evidence for specific effects of periodontal treatment procedures on increasing ischaemic cardiovascular risk. Two observational

studies reported no effect of “*invasive dental treatment*” in elevating ischaemic cardiovascular risk (Chen et al., 2019; Nordendahl et al., 2018), and one study suggested a minimal increased risk within 4 weeks following treatment (Minassian, D’Aiuto, Hingorani, & Smeeth, 2010).

Chen et al. (2019) performed a case-crossover and self-controlled case series using the Taiwanese National Health Insurance Research Database, including over 110,000 Myocardial Infarction cases and 290,000 ischaemic stroke patients between 1999 and 2014. They reported a non-significant increase in the incidence of myocardial infarction within the first 24 weeks following “*invasive dental treatment*” (including periodontal procedures) except for a modest risk of myocardial infarction during the first week for patients without other comorbidities (OR = 1.31, 95% CI [1.08; 1.58], after 3 days).

A registry-based case-control study between 2011 and 2013 including 51,880 cases who underwent an “*invasive dental procedure*” compared to 246,978 controls reported no association with an increased incidence of myocardial infarction (OR 0.98, 95% CI [0.91; 1.06]) (Nordendahl et al., 2018).

Minassian et al. (2010) performed a self-controlled case series including nearly 10 million participants included in an insurance database from 2002 and 2006 in the United States. The analysis showed that *invasive dental treatment* (largely comprising of tooth extractions and only 4% being non-surgical and surgical periodontal procedures) is associated with an increased risk of incident acute cardiovascular events (IR = 1.5, 95% CI [1.09; 2.06]) within the first 4 weeks of treatment recorded.

In summary, the Group concluded that delivering periodontal treatment is safe with regard to cardiovascular risk.

5.1.2 | In patients with established CVD

There is limited evidence on the effects of “*invasive dental treatment*” on the incidence of ischaemic events in patients with established CVD or after an event.

A small RCT on the effects of the treatment of periodontitis on CVD biomarkers in patients with established CVD (Montenegro et al., 2019) showed no cardiovascular adverse events within 3 months of completion of scaling and root planing (periodontal therapy).

In the PAVE feasibility randomized secondary prevention trial, provision of periodontal scaling and root planing treatment in patients with established CVD did not increase the incidence of cardiovascular events compared to the control group (community treatment) within 6 months (Beck et al., 2008).

In summary, the Group concluded that delivering periodontal treatment is safe with regard to cardiovascular risk in patients with established CVD.

5.2 | What is the perioperative bleeding risk when performing periodontal therapy?

Periodontal treatment consists of numerous procedures with different levels of bleeding risk. This risk of bleeding is however low in the

vast majority of procedures, and it can be easily controlled with local haemostatic measures.

Perioperative bleeding risk varies according to the extent and invasiveness of the periodontal procedure performed. The majority of periodontal procedures may be grouped within the ESC/AHA/EHRA (Steffel et al., 2018a, 2018b). Low bleeding risk group (frequency less than 1% of post-operative bleeding) group: supragingival polishing, non-surgical periodontal treatment, conventional surgical periodontal treatment (conservative, resective or regenerative), tooth extractions and dental implant placement. Moderate bleeding risk (frequency between 2 and 5%) may be observed in major autogenous bone augmentation procedures such as block bone harvesting, sinus floor elevation and procedures where healing is by secondary intention, such as free gingival grafting. Appendix S1 summarized the main recommendations for patients with antithrombotic therapy when performing periodontal therapy.

5.2.1 | In patients undergoing antiplatelet therapy

Individuals undergoing single acetylsalicylic acid (ASA) therapy (aspirin) in different therapeutic dosages, as well as therapy with clopidogrel, ticlopidine or ticagrelor, show no statistically significant differences in frequency of bleeding events when compared to controls, that is subjects not undergoing antiplatelet therapy (Doganay, Atalay, Karadag, Aga, & Tugrul, 2018; Lillis, Ziakas, Koskinas, Tsirlis, & Giannoglou, 2011).

Dual antiplatelet therapy, most commonly ASA in combination with clopidogrel, may pose a certain risk for post-operative bleeding complications; however, it appears that these haemorrhagic events may be managed safely with local haemostatic measures (Napenas et al., 2009; Nathwani & Martin, 2016).

Thus, current evidence does not support discontinuation of antiplatelet therapy before dental procedures, irrespective of the type of therapy employed (single or dual antiplatelet therapy) or the type of procedure performed (single, multiple tooth extractions, non-surgical and surgical periodontal therapy and dental implant procedures).

5.2.2 | In patients undergoing anticoagulant therapy

Vitamin K antagonists

In patients taking oral anticoagulant therapy (vitamin K antagonists, VKA) and undergoing dental extraction, minor dental procedures and dental implant placement do not seem to increase the risk of bleeding compared to patients who discontinue oral anticoagulant therapy (Shi, Xu, Zhang, Zhang, & Liu, 2017; Yang, Shi, Liu, Li, & Xu, 2016). There may be a higher post-operative bleeding risk in patients continuing VKA and undergoing either minor dental surgery or other higher-risk procedures when compared to non-VKA patients (Biedermann et al., 2017; Shi et al., 2017), but local haemostatic agents appear to be effective in controlling post-operative bleeding (Madrid & Sanz, 2009).

Novel/direct anticoagulants (DOAC/NOAC)

Limited trials and evidence are available on the management of patients on novel oral anticoagulant (NOAC) therapy undergoing dental treatment; hence, the Group concluded that further studies regarding dental procedures in these patients are strongly encouraged.

It appears there is no need for interruption of NOAC therapy in most dental treatments, due to a low incidence of bleeding complications, which can be successfully managed with local haemostatic measures when comparing groups continuing NOAC and groups discontinuing NOAC therapy (Kwak et al., 2019; Lababidi et al., 2018; Patel et al., 2017; Yagyu et al., 2017) and with reported timing of discontinuation and reinstitution varying greatly. When comparing NOAC patients with healthy individuals, there seems to be a higher incidence of delayed bleeding (2 days and later) in those patients who do not discontinue NOAC therapy (Miclotte et al., 2017).

6 | RECOMMENDATIONS

6.1 | Recommendations for oral health professionals for use in dental practice/office for people with cardiovascular disease (CVD)

- Patients with periodontitis should be advised that there is a higher risk for cardiovascular diseases, such as myocardial infarction or stroke, and as such, they should actively manage all their cardiovascular risk factors (smoking, exercise, excess weight, blood pressure, lipid and glucose management, and sufficient periodontal therapy and periodontal maintenance).
- Patients with periodontitis and a diagnosis of CVD should be informed that they may be at higher risk for subsequent CVD complications, and therefore, they should regularly adhere to the recommended dental therapeutic, maintenance and preventive regimes.
- Patients collect a careful history to assess for CVD risk factors, such as diabetes, obesity, smoking, hypertension, hyperlipidaemia and hyperglycaemia. Patients suggest that the patient consults his/her physician if any of these risk factors are not appropriately controlled.
- Oral health education should be provided to all patients with periodontitis and a tailored oral hygiene regime, including twice-daily brushing, interdental cleaning and, in some cases, the use of adjunctive chemical plaque control, may be appropriate.
- People presenting with a diagnosis of CVD should receive a thorough oral examination, which embeds a comprehensive periodontal evaluation, including full-mouth probing and bleeding scores.
- If no periodontitis is diagnosed initially, patients with CVD should be placed on a preventive care regime and monitored regularly (at least once a year) for changes in periodontal status.
- In people with CVD, if periodontitis is diagnosed, they should be

managed as soon as their cardiovascular status permits.

- o Irrespective of the level of CVD or specific medication, non-surgical periodontal therapy should be provided, preferably in several 30- to 45-min sessions, in order to minimize a spike of acute systemic inflammation
- o Surgical periodontal and implant therapy when indicated should be provided in a similar manner as in patients without CVD.

However, attention should be paid to:

- Hypertension. It is recommended to measure the patients' blood pressure (after appropriate relaxation) before the surgical intervention, and in cases of high blood pressure (above 180/100 according to expert opinion), the surgery should be postponed until the patient's blood pressure is stabilized.
- Medication with antiplatelet and anticoagulant drugs. Since periodontal and implant surgical procedures usually impart only a low-to-medium risk of bleeding in general terms, the dentist should not change a patient's medication, or in cases of doubt, he/she should consult the physician/cardiologist prior to the surgical intervention. Consideration should also be given to the local management of bleeding complications that may arise.

Current AHA/ACC/SCAI/ACS/ADA/ESC/ACCP guidelines on perioperative management of antithrombotic therapy do not suggest discontinuation of anti-platelet therapy for low bleeding risk procedures (Douketis et al., 2012; Grines, Bonow, & Casey, 2007; Kristensen et al., 2014).

Various approaches for peri-operative management of anti-coagulant therapy have been suggested. The Group reviewed the guidelines on perioperative management of vitamin K antagonists (VKA) and suggested discontinuation of medication treatment if the INR is 4 or below for low or medium bleeding risk procedures (Perry, Noakes, Helliwell, & British Dental, 2007). However, if the INR (internationalized normalized ratio) is 3.5 or above, the expert group recommends that dental clinicians seek advice and consult with the responsible medical professional. Management of high thrombo-embolic risk cases should be collaborative in consultation with the medical professional responsible for VKA therapy (Kristensen et al., 2014; Valgimigli et al., 2018).

After reviewing novel anticoagulant (non-VKA) and direct anti-coagulant (NOAC/DOAC) therapies guidelines, the Group concluded that for low bleeding risk periodontal procedures no discontinuation of anticoagulants is recommended (Steffel et al., 2018a, 2018b). These procedures could be performed 18-24 hrs after the last intake (depending on a renal function assessment for the medication in question) and then restart 6 hrs following treatment. The expert group, however, strongly recommends that the dental clinician should consult with the responsible medical professional. When a medium bleeding risk periodontal procedure is planned, discontinuation of therapy should be agreed with the medical professional responsible for and/or prescribing the anticoagulant therapy.

Lastly, in cases of combined antiplatelet and anticoagulant therapies that pertain patients with the highest thrombotic and ischaemic risk (i.e. chronic atrial fibrillation or after an acute myocardial infarction or recent coronary stenting), when periodontal procedures (either of low or medium bleeding risk) are required, any alterations in medication should be discussed and agreed upon with the responsible medical professional (Steffel et al., 2018a, 2018b). In elective periodontal procedures, the operation should be delayed until after treatment stabilization and appropriate consultation with the medical specialist.

In cases of triple therapy (dual antiplatelet and one anticoagulant) or one anticoagulant plus one antiplatelet, such patients need individualized management by the responsible medical professional according to their thrombotic and haemorrhagic risk (Valgimigli et al., 2018).

It is important to highlight that local haemostatic agents (such as oxidized cellulose, absorbable gelatin sponges, sutures, tranexamic acid mouthwashes, compressive gauze soaked in tranexamic acid) should be used and dental clinicians should consider the confounding effect of local anaesthetic with vasoconstrictors.

- Patients with a risk of endocarditis should be premedicated with antibiotics following current guidelines (such as the European or the American guidelines).
- People with cardiovascular disease who have extensive tooth loss should be encouraged to pursue dental rehabilitation to restore adequate mastication for proper nutrition.

People without a diagnosis of CVD, but with risk factors for CVD should be informed about their CVD risk and referred to a physician for appropriate risk assessment, diagnostic testing and follow-up care. For oral health professionals, risk assessment may be performed based upon the recommendations of the *European Society of Cardiology (Systematic COronary Risk Evaluation, SCORE)* (Sixth Joint Task Force of the European Society of Cardiology & Other Societies on Cardiovascular Disease Prevention in Clinical Practice, 2016).

6.2 | Recommendations for physicians and other medical health professions for use in cardiology practice

Because of the potential negative impact of periodontitis on CVD complications, the following recommendations are made:

- Patients with CVD should be advised that periodontitis may have a negative impact on CVD and may also increase the risk of CVD events.
- Patients should be advised that effective periodontal therapy may have a positive impact upon CV health.
- For people with CVD, physicians should ask about a prior diagnosis of periodontitis. If a positive diagnosis has been made, the physician should seek to ascertain that appropriate periodontal care and maintenance are being provided.
- Patients with CVD should be asked about any signs and symptoms of periodontitis, including bleeding gums during brushing or

eating, loose teeth, spacing or spreading/drift of the teeth, oral malodor and/or abscesses of the gums or gingival suppuration.

- o If a positive history is elicited, then a prompt periodontal evaluation should be recommended before their scheduled annual check-up.
- o In the case of a negative history, people with CVD should be advised to check for the above symptoms, and if a positive sign appears, they should visit their dentist at least once per year.
- For all patients with newly diagnosed CVD, referral for a periodontal examination should occur as part of their ongoing management of CVD. Even if no periodontitis is diagnosed initially, an annual oral/dental check-up is recommended.
- The physician should liaise with the dental surgeon over periodontitis management in CVD patients on anticoagulant/antiplatelet therapy prior to the oral intervention and/or periodontal surgery, to avoid excess bleeding or the risk of ischaemic events.

6.3 | Recommendations for patients at the dental surgery/ office who have CVD or are found to be at risk of CVD

- People with CVD must be aware that gum disease is a chronic condition, which may aggravate their CVD and requires lifelong attention and professional care.
- There is a need to clean the teeth and gums very carefully at home. Personalized advice will be provided by the oral health professional.

This may include the following:

- Twice-daily brushing with either a manual or electric toothbrush;
- Cleaning between teeth using interdental brushes where they fit, and where they do not fit, then flossing may be useful;
- Use of specific dentifrices and/or mouth rinses with proven activity against dental plaque, if advised by oral health professionals;
- If left untreated, gum disease can lead to tooth loss and may also make CVD preventive measures harder to control;
- Gum disease may be present and deteriorate with no apparent symptoms, so the dentist should advise their patient that even without current gum disease, they should still receive regular dental check-ups as part of managing their CVD.

Dentists should be able to identify the early signs of gum disease, but patients should also suspect gum disease if noticing:

- Red or swollen gums;
- Bleeding from the gums or blood in the sink after toothbrushing;
- Foul taste;
- Longer looking teeth;
- Loose teeth;
- Increasing spaces between teeth/ teeth moving apart;
- Calculus (tartar) on teeth.

Patients should inform their dentist about the outcome of their visits to the physician and provide an update on their CVD history and any changes in medications. Patients should inform the dentist if they are on anticoagulant therapy.

Patients should understand that it is important to keep their mouth and whole body as healthy as possible with regular dental and medical visits.

6.4 | Recommendations for patients with CVD at the physician's practice/office

6.4.1 | Why should I have my gums checked?

If your physician has told you that you have cardiovascular disease (CVD), you should make an appointment with a dental surgeon to have your mouth and gums checked.

This is because people with CVD may have a higher chance of getting further complications when they have gum disease. The earlier you seek help, the better the outcome will be.

6.4.2 | What should I look for that may tell me I have problems with my gums?

You may have gum disease if you have ever noticed:

- Red or swollen gums;
- Bleeding from your gums or blood in the sink after you brush your teeth;
- Foul taste;
- Longer looking teeth;
- Loose teeth;
- Increasing spaces between your teeth, or your teeth drifting apart;
- Calculus (tartar) on your teeth.

If you have noticed any of these problems, it is important to see a dentist as soon as possible.

6.4.3 | Can I have gum disease without these signs being present?

Gum disease may also be present and get worse with no apparent signs to you that you have it, especially if you smoke, so even if you do not think you have gum disease now, you should still have annual check-ups of your mouth as part of managing your CVD. Your dentist will be able to pick up early signs of gum disease.

6.4.4 | What can I do to prevent gum disease?

You need to clean your teeth and gums twice daily at home for a minimum of 2 min. Also, cleaning between your teeth daily is important and your oral health professional will show you how to do this. You should visit a dental surgeon as soon as possible for

a diagnosis and advice on what you need to do. It is important to keep your mouth as healthy as possible with regular oral and dental care, according to the recommendations of your oral health professional.

ORCID

Mariano Sanz  <https://orcid.org/0000-0002-6293-5755>

Søren Jepsen  <https://orcid.org/0000-0002-4160-5837>

Filippo Graziani  <https://orcid.org/0000-0001-8780-7306>

David Herrera  <https://orcid.org/0000-0002-5554-2777>

Bruno Loos  <https://orcid.org/0000-0002-8794-552X>

Phoebus Madianos  <https://orcid.org/0000-0002-5990-5723>

Lior Shapira  <https://orcid.org/0000-0001-9145-5155>

Maurizio Tonetti  <https://orcid.org/0000-0002-2743-0137>

REFERENCES

- Aarabi, G., Zeller, T., Seedorf, H., Reissmann, D. R., Heydecke, G., Schaefer, A. S., & Seedorf, U. (2017). Genetic susceptibility contributing to periodontal and cardiovascular disease. *Journal of Dental Research*, 96, 610–617. <https://doi.org/10.1177/0022034517699786>
- Ahn, Y. B., Shin, M. S., Han, D. H., Sukhbaatar, M., Kim, M. S., Shin, H. S., & Kim, H. D. (2016). Periodontitis is associated with the risk of subclinical atherosclerosis and peripheral arterial disease in Korean adults. *Atherosclerosis*, 311, e318. <https://doi.org/10.1016/j.atherosclerosis.2016.07.898>
- Armingohar, Z., Jørgensen, J. J., Kristoffersen, A. K., Abesha-Belay, E., & Olsen, I. (2014). Bacteria and bacterial DNA in atherosclerotic plaque and aneurysmal wall biopsies from patients with and without periodontitis. *Journal of Oral Microbiology*, 6. <https://doi.org/10.3402/jom.v6.23408>
- Arvanitidis, E., Bizzarro, S., Alvarez Rodriguez, E., Loos, B. G., & Nicu, E. A. (2017). Reduced platelet hyper-reactivity and platelet-leukocyte aggregation after periodontal therapy. *Thrombosis Journal*, 15, 5. <https://doi.org/10.1186/s12959-016-0125-x>
- Balejo, R. D. P., Cortelli, J. R., Costa, F. O., Cyrino, R. M., Aquino, D. R., Cogo-Müller, K., ... Cortelli, S. C. (2017). Effects of chlorhexidine pre-procedural rinse on bacteremia in periodontal patients: A randomized clinical trial. *Journal of Applied Oral Science*, 25, 586–595. <https://doi.org/10.1590/1678-7757-2017-0112>
- Beck, J. D., Couper, D. J., Falkner, K. L., Graham, S. P., Grossi, S. G., Gunsolley, J. C., ... Genco, R. J. (2008). The Periodontitis and Vascular Events (PAVE) pilot study: Adverse events. *Journal of Periodontology*, 79, 90–96. <https://doi.org/10.1902/jop.2008.070223>
- Belanger, M., Kozarov, E., Song, H., Whitlock, J., & Progulske-Fox, A. (2012). Both the unique and repeat regions of the *Porphyromonas gingivalis* hemagglutinin A are involved in adhesion and invasion of host cells. *Anaerobe*, 18, 128–134. <https://doi.org/10.1016/j.anaerobe.2011.10.005>
- Bertl, K., Parllaku, A., Pandis, N., Buhlin, K., Klinge, B., & Stavropoulos, A. (2017). The effect of local and systemic statin use as an adjunct to non-surgical and surgical periodontal therapy-A systematic review and meta-analysis. *Journal of Dentistry*, 67, 18–28. <https://doi.org/10.1016/j.jdent.2017.08.011>
- Bertl, K., Steiner, I., Pandis, N., Buhlin, K., Klinge, B., & Stavropoulos, A. (2018). Statins in nonsurgical and surgical periodontal therapy.

- A systematic review and meta-analysis of preclinical in vivo trials. *Journal of Periodontal Research*, 53, 267–287. <https://doi.org/10.1111/jre.12514>
- Biedermann, J. S., Rademacher, W. M. H., Hazendonk, H. C. A. M., van Diermen, D. E., Leebeek, F. W. G., Rozema, F. R., & Kruip, M. J. H. A. (2017). Predictors of oral cavity bleeding and clinical outcome after dental procedures in patients on vitamin K antagonists. *Thrombosis and Haemostasis*, 117, 1432–1439. <https://doi.org/10.1160/TH17-01-0040>
- Boillot, A., Range, H., Danchin, N., Kotti, S., Cosler, G., Czernichow, S., ... Simon, T. (2016). Periodontopathogens antibodies and major adverse events following an acute myocardial infarction: Results from the French Registry of Acute ST-Elevation and Non-ST-Elevation Myocardial Infarction (FAST-MI). *Journal of Epidemiology and Community Health*, 70, 1236–1241. <https://doi.org/10.1136/jech-2015-207043>
- Caula, A. L., Lira-Junior, R., Tinoco, E. M., & Fischer, R. G. (2014). The effect of periodontal therapy on cardiovascular risk markers: A 6-month randomized clinical trial. *Journal of Clinical Periodontology*, 41, 875–882. <https://doi.org/10.1111/jcpe.12290>
- Centers for Disease Control and Prevention (2011). *Healthy aging at a glance 2011*. Atlanta, GA: Centers for Disease Control and Prevention.
- Centers for Disease Control and Prevention (2013) *The state of aging and health in America 2013*. Atlanta, GA: Centers for Disease Control and Prevention, US Dept of Health and Human Services.
- Chandy, S., Joseph, K., Sankaranarayanan, A., Issac, A., Babu, G., Wilson, B., & Joseph, J. (2017). Evaluation of C-reactive protein and fibrinogen in patients with chronic and aggressive periodontitis: A clinico-biochemical study. *Journal of Clinical and Diagnostic Research*, 11, Zc41–zc45. <https://doi.org/10.7860/jcdr/2017/23100.9552>
- Chapple, I. L., Brock, G. R., Milward, M. R., Ling, N., & Matthews, J. B. (2007). Compromised GCF total antioxidant capacity in periodontitis: Cause or effect? *Journal of Clinical Periodontology*, 34, 103–110. <https://doi.org/10.1111/j.1600-051X.2006.01029.x>
- Chapple, I. L., Genco, R., & Working Group 2 of the Joint EFP/AAP Workshop (2013). Diabetes and periodontal diseases: Consensus report of the Joint EFP/AAP Workshop on Periodontitis and Systemic Diseases. *Journal of Clinical Periodontology*, 40(Suppl 14), S106–112. <https://doi.org/10.1111/jcpe.12077>
- Chen, D. Y., Lin, C. H., Chen, Y. M., & Chen, H. H. (2016). Risk of atrial fibrillation or flutter associated with periodontitis: A nationwide, population-based cohort study. *PLoS ONE*, 11, e0165601. <https://doi.org/10.1371/journal.pone.0165601>
- Chen, T. T., D'Aiuto, F., Yeh, Y. C., Lai, M. S., Chien, K. L., & Tu, Y. K. (2019). Risk of myocardial infarction and ischemic stroke after dental treatments. *Journal of Dental Research*, 98, 157–163. <https://doi.org/10.1177/0022034518805745>
- Chukkapalli, S. S., Velsko, I. M., Rivera-Kweh, M. F., Zheng, D., Lucas, A. R., & Kesavalu, L. (2015). Polymicrobial oral infection with four periodontal bacteria orchestrates a distinct inflammatory response and atherosclerosis in ApoE null mice. *PLoS ONE*, 10, e0143291. <https://doi.org/10.1371/journal.pone.0143291>
- Couper, D. J., Beck, J. D., Falkner, K. L., Graham, S. P., Grossi, S. G., Gunsolley, J. C., ... Genco, R. J. (2008). The Periodontitis and Vascular Events (PAVE) pilot study: Recruitment, retention, and community care controls. *Journal of Periodontology*, 79, 80–89. <https://doi.org/10.1902/jop.2008.070216>
- D'Aiuto, F., Gkraniyas, N., Bhowruth, D., Khan, T., Orlandi, M., Suvan, J., ... TASTE Group (2018). Systemic effects of periodontitis treatment in patients with type 2 diabetes: A 12 month, single-centre, investigator-masked, randomised trial. *Lancet Diabetes and Endocrinology*, 6, 954–965. [https://doi.org/10.1016/S2213-8587\(18\)30038-X](https://doi.org/10.1016/S2213-8587(18)30038-X)
- D'Aiuto, F., Orlandi, M., & Gunsolley, J. C. (2013). Evidence that periodontal treatment improves biomarkers and CVD outcomes. *Journal of Clinical Periodontology*, 40(Suppl 14), S85–105. <https://doi.org/10.1111/jcpe.12061>
- de Boer, S. P., Cheng, J. M., Range, H., Garcia-Garcia, H. M., Heo, J. H., Akkerhuis, K. M., ... Kardys, I. (2014). Antibodies to periodontal pathogens are associated with coronary plaque remodeling but not with vulnerability or burden. *Atherosclerosis*, 237, 84–91. <https://doi.org/10.1016/j.atherosclerosis.2014.08.050>
- de Oliveira, C., Watt, R., & Hamer, M. (2010). Toothbrushing, inflammation, and risk of cardiovascular disease: Results from Scottish Health Survey. *British Medical Journal*, 340, c2451. <https://doi.org/10.1136/bmj.c2451>
- Deepti, Tewari, S., Narula, S. C., Singhal, S. R., & Sharma, R. K. (2017). Effect of non-surgical periodontal therapy along with myo-inositol on high-sensitivity C-reactive protein and insulin resistance in women with polycystic ovary syndrome and chronic periodontitis: A randomized controlled trial. *Journal of Periodontology*, 88, 999–1011. <https://doi.org/10.1902/jop.2017.170121>
- Demmer, R. T., Trinquart, L., Zuk, A., Fu, B. C., Blomkvist, J., Michalowicz, B. S., ... Desvarieux, M. (2013). The influence of anti-infective periodontal treatment on C-reactive protein: A systematic review and meta-analysis of randomized controlled trials. *PLoS ONE*, 8, e77441. <https://doi.org/10.1371/journal.pone.0077441>
- Dietrich, T., Sharma, P., Walter, C., Weston, P., & Beck, J. (2013). The epidemiological evidence behind the association between periodontitis and incident atherosclerotic cardiovascular disease. *Journal of Clinical Periodontology*, 40, S70–84. <https://doi.org/10.1111/jcpe.12062>
- Divaris, K., Monda, K. L., North, K. E., Olshan, A. F., Lange, E. M., Moss, K., ... Offenbacher, S. (2012). Genome-wide association study of periodontal pathogen colonization. *Journal of Dental Research*, 91, 21S–28S. <https://doi.org/10.1177/0022034512447951>
- Doganay, O., Atalay, B., Karadag, E., Aga, U., & Tugrul, M. (2018). Bleeding frequency of patients taking ticagrelor, aspirin, clopidogrel, and dual antiplatelet therapy after tooth extraction and minor oral surgery. *Journal of the American Dental Association*, 149, 132–138. <https://doi.org/10.1016/j.adaj.2017.09.052>
- Dorn, J. M., Genco, R. J., Grossi, S. G., Falkner, K. L., Hovey, K. M., Iacoviello, L., & Trevisan, M. (2010). Periodontal disease and recurrent cardiovascular events in survivors of myocardial infarction (MI): The Western New York Acute MI Study. *Journal of Periodontology*, 81, 502–511. <https://doi.org/10.1902/jop.2009.090499>
- Douketis, J. D., Spyropoulos, A. C., Spencer, F. A., Mayr, M., Jaffer, A. K., Eckman, M. H., ... Kunz, R. (2012). Perioperative management of antithrombotic therapy: Antithrombotic Therapy and Prevention of Thrombosis, 9th ed: American College of Chest Physicians Evidence-Based Clinical Practice Guidelines. *Chest*, 141, e326S–e350S. <https://doi.org/10.1378/chest.11-2298>
- Ernst, F. D., Uhr, K., Teumer, A., Fanghanel, J., Schulz, S., Noack, B., ... Kocher, T. (2010). Replication of the association of chromosomal region 9p21.3 with generalized aggressive periodontitis (gAgP) using an independent case-control cohort. *BMC Medical Genetics*, 11, 119. <https://doi.org/10.1186/1471-2350-11-119>
- Estanislau, I. M., Terceiro, I. R., Lisboa, M. R., Teles Pde, B., Carvalho Rde, S., Martins, R. S., & Moreira, M. M. (2015). Pleiotropic effects of statins on the treatment of chronic periodontitis—a systematic review. *British Journal of Clinical Pharmacology*, 79, 877–885. <https://doi.org/10.1111/bcp.12564>
- Fajardo, M. E., Rocha, M. L., Sanchez-Marin, F. J., & Espinosa-Chavez, E. J. (2010). Effect of atorvastatin on chronic periodontitis: A randomized pilot study. *Journal of Clinical Periodontology*, 37, 1016–1022. <https://doi.org/10.1111/j.1600-051X.2010.01619.x>
- Fentoglu, O., Kirzioglu, F. Y., Ozdem, M., Kocak, H., Sutcu, R., & Sert, T. (2012). Proinflammatory cytokine levels in hyperlipidemic patients with periodontitis after periodontal treatment. *Oral Diseases*, 18, 299–306. <https://doi.org/10.1111/j.1601-0825.2011.01880.x>

- Freitas, C. O., Gomes-Filho, I. S., Naves, R. C., Nogueira Filho, G. R., Cruz, S. S., Santos, C. A., ... Barbosa, M. D. (2012). Influence of periodontal therapy on C-reactive protein level: A systematic review and meta-analysis. *Journal of Applied Oral Science*, 20, 1–8. <https://doi.org/10.1590/S1678-77572012000100002>
- Fu, Y. W., Li, X. X., Xu, H. Z., Gong, Y. Q., & Yang, Y. (2016). Effects of periodontal therapy on serum lipid profile and proinflammatory cytokines in patients with hyperlipidemia: A randomized controlled trial. *Clinical Oral Investigations*, 20, 1263–1269. <https://doi.org/10.1007/s00784-015-1621-2>
- G. B. D. Disease Injury and Incidence and Prevalence Collaborators (2018). Global, regional, and national incidence, prevalence, and years lived with disability for 354 diseases and injuries for 195 countries and territories, 1990–2017: A systematic analysis for the Global Burden of Disease Study 2017. *Lancet*, 392, 1789–1858. [https://doi.org/10.1016/S0140-6736\(18\)32279-7](https://doi.org/10.1016/S0140-6736(18)32279-7)
- G. B. D. Risk Factors Collaborators (2016). Global, regional, and national comparative risk assessment of 79 behavioural, environmental and occupational, and metabolic risks or clusters of risks, 1990–2015: A systematic analysis for the Global Burden of Disease Study 2015. *Lancet*, 388, 1659–1724. [https://doi.org/10.1016/S0140-6736\(16\)31679-8](https://doi.org/10.1016/S0140-6736(16)31679-8)
- Garrett, I. R., Gutierrez, G., & Mundy, G. R. (2001). Statins and bone formation. *Current Pharmaceutical Design*, 7, 715–736.
- Graziani, F., Cei, S., Orlandi, M., Gennai, S., Gabriele, M., Filice, N., ... D'Aiuto, F. (2015). Acute-phase response following full-mouth versus quadrant non-surgical periodontal treatment: A randomized clinical trial. *Journal of Clinical Periodontology*, 42, 843–852. <https://doi.org/10.1111/jcpe.12451>
- Grines, C. L., Bonow, R. O., Casey, D. E. Jr, Gardner, T. J., Lockhart, P. B., Moliterno, D. J., ... American College of Physicians (2007). Prevention of premature discontinuation of dual antiplatelet therapy in patients with coronary artery stents: A science advisory from the American Heart Association, American College of Cardiology, Society for Cardiovascular Angiography and Interventions, American College of Surgeons, and American Dental Association, with representation from the American College of Physicians. *Journal of the American Dental Association*, 138, 652–655. <https://doi.org/10.14219/jada.archive.2007.0237>
- Hada, D. S., Garg, S., Ramteke, G. B., & Rathe, M. S. (2015). Effect of non-surgical periodontal treatment on clinical and biochemical risk markers of cardiovascular disease: A randomized trial. *Journal of Periodontology*, 86, 1201–1211. <https://doi.org/10.1902/jop.2015.150249>
- Herrera, D., Molina, A., Buhlin, K., & Klinge, B. (2019) Periodontal diseases and association with atherosclerotic disease. *Periodontology* 2000, (in press).
- Holdt, M. H., & Teupser, D. (2015). Genetic background of atherosclerosis and its risk factors. In *The ESC textbook of preventive cardiology* (pp. 21–25). Oxford, UK: Oxford University Press.
- Holmlund, A., Lampa, E., & Lind, L. (2017). Poor response to periodontal treatment may predict future cardiovascular disease. *Journal of Dental Research*, 96, 768–773. <https://doi.org/10.1177/0022034517701901>
- Hubberten, M., Bochenek, G., Chen, H., Hasler, R., Wiehe, R., Rosenstiel, P., ... Schaefer, A. S. (2019). Linear isoforms of the long noncoding RNA CDKN2B-AS1 regulate the c-myc-enhancer binding factor RBMS1. *European Journal of Human Genetics*, 27, 80–89. <https://doi.org/10.1038/s41431-018-0210-7>
- Ioannidou, E., Malekzadeh, T., & Dongari-Bagtzoglou, A. (2006). Effect of periodontal treatment on serum C-reactive protein levels: A systematic review and meta-analysis. *Journal of Periodontology*, 77, 1635–1642. <https://doi.org/10.1902/jop.2006.050443>
- Joseph, P., Leong, D., McKee, M., Anand, S. S., Schwalm, J.-D., Teo, K., ... Yusuf, S. (2017). Reducing the global burden of cardiovascular disease. Part 1: The epidemiology and risk factors. *Circulatory Research*, 121(6), 677–694. <https://doi.org/10.1161/CIRCRES.ESAHA.117.308903>
- Kapellas, K., Maple-Brown, L. J., Jamieson, L. M., Do, L. G., O'Dea, K., Brown, A., ... Skilton, M. R. (2014). Effect of periodontal therapy on arterial structure and function among aboriginal Australians: A randomized, controlled trial. *Hypertension*, 64, 702–708. <https://doi.org/10.1161/HYPERTENSIONAHA.114.03359>
- Kassebaum, N. J., Bernabe, E., Dahiya, M., Bhandari, B., Murray, C. J., & Marcenes, W. (2014). Global burden of severe periodontitis in 1990–2010: A systematic review and meta-regression. *Journal of Dental Research*, 93, 1045–1053. <https://doi.org/10.1177/0022034514552491>
- Kassebaum, N. J., Smith, A. G. C., Bernabé, E., Fleming, T. D., Reynolds, A. E., Vos, T., ... Oral Health Collaborators (2017). Global, regional, and national prevalence, incidence, and disability adjusted life years for oral conditions for 195 countries, 1990–2015: A systematic analysis for the global burden of diseases, injuries, and risk factors. *Journal of Dental Research*, 96(4), 380–387. <https://doi.org/10.1177/0022034517693566>
- Kaushal, S., Singh, A. K., Lal, N., Das, S. K., & Mahdi, A. A. (2019). Effect of periodontal therapy on disease activity in patients of rheumatoid arthritis with chronic periodontitis. *Journal of Oral Biology and Craniofacial Research*, 9, 128–132. <https://doi.org/10.1016/j.jobcr.2019.02.002>
- Koh, K. K., Son, J. W., Ahn, J. Y., Jin, D. K., Kim, H. S., Choi, Y. M., ... Shin, E. K. (2002). Comparative effects of diet and statin on NO bioactivity and matrix metalloproteinases in hypercholesterolemic patients with coronary artery disease. *Arteriosclerosis, Thrombosis, and Vascular Biology*, 22, e19–23. <https://doi.org/10.1161/01.ATV.0000030997.02059.BB>
- Koppolu, P., Durvasula, S., Palaparthi, R., Rao, M., Sagar, V., Reddy, S. K., & Lingam, S. (2013). Estimate of CRP and TNF-alpha level before and after periodontal therapy in cardiovascular disease patients. *Pan African Medical Journal*, 15, 92. <https://doi.org/10.11604/pamj.2013.15.92.2326>
- Kozarov, E. V., Dorn, B. R., Shelburne, C. E., Dunn, W. A. Jr, & Progluske-Fox, A. (2005). Human atherosclerotic plaque contains viable invasive *Actinobacillus actinomycetemcomitans* and *Porphyromonas gingivalis*. *Arteriosclerosis, Thrombosis, and Vascular Biology*, 25, e17–18. <https://doi.org/10.1161/01.ATV.0000155018.67835.1a>
- Kristensen, S. D., Knuuti, J., Saraste, A., Anker, S., Botker, H. E., Hert, S. D., ... Authors/Task Force Members (2014). 2014 ESC/ESA Guidelines on non-cardiac surgery: Cardiovascular assessment and management: The Joint Task Force on non-cardiac surgery: Cardiovascular assessment and management of the European Society of Cardiology (ESC) and the European Society of Anaesthesiology (ESA). *European Heart Journal*, 35, 2383–2431. <https://doi.org/10.1093/eurheartj/ehu282>
- Kwak, E. J., Nam, S., Park, K. M., Kim, S. Y., Huh, J., & Park, W. (2019). Bleeding related to dental treatment in patients taking novel oral anti-coagulants (NOACs): A retrospective study. *Clinical Oral Investigations*, 23, 477–484. <https://doi.org/10.1007/s00784-018-2458-2>
- Lababidi, E., Breik, O., Savage, J., Engelbrecht, H., Kumar, R., & Crossley, C. W. (2018). Assessing an oral surgery specific protocol for patients on direct oral anticoagulants: A retrospective controlled cohort study. *International Journal of Oral and Maxillofacial Surgery*, 47, 940–946. <https://doi.org/10.1016/j.ijom.2018.03.009>
- Lee, Y. L., Hu, H. Y., Chou, P., & Chu, D. (2015). Dental prophylaxis decreases the risk of acute myocardial infarction: A nationwide population-based study in Taiwan. *Clinical Interventions in Aging*, 10, 175–182. <https://doi.org/10.2147/CIA.S67854>
- Li, C., Lv, Z., Shi, Z., Zhu, Y., Wu, Y., Li, L., & Iheozor-Ejiofor, Z. (2017). Periodontal therapy for the management of cardiovascular disease in patients with chronic periodontitis. *Cochrane Database of Systematic*

- Reviews, 11, CD009197. <https://doi.org/10.1002/14651858.CD009197.pub3>
- Lillis, T., Ziakas, A., Koskinas, K., Tsiolis, A., & Giannoglou, G. (2011). Safety of dental extractions during uninterrupted single or dual antiplatelet treatment. *American Journal of Cardiology*, 108, 964–967. <https://doi.org/10.1016/j.amjcard.2011.05.029>
- Linden, G. J., Linden, K., Yarnell, J., Evans, A., Kee, F., & Patterson, C. C. (2012). All-cause mortality and periodontitis in 60–70-year-old men: A prospective cohort study. *Journal of Clinical Periodontology*, 39, 940–946. <https://doi.org/10.1111/j.1600-051X.2012.01923.x>
- Linden, G. J., Lyons, A., & Scannapieco, F. A. (2013). Periodontal systemic associations: Review of the evidence. *Journal of Clinical Periodontology*, 40(Suppl 14), S8–19. <https://doi.org/10.1111/jcpe.12064>
- Lindy, O., Suomalainen, K., Mäkelä, M., & Lindy, S. (2008). Statin use is associated with fewer periodontal lesions: A retrospective study. *BMC Oral Health*, 15, 16. <https://doi.org/10.1186/1472-6831-8-16>
- Ling, M. R., Chapple, I. L., & Matthews, J. B. (2016). Neutrophil superoxide release and plasma C-reactive protein levels pre- and post-periodontal therapy. *Journal of Clinical Periodontology*, 43, 652–658. <https://doi.org/10.1111/jcpe.12575>
- Liu, S., Bertl, K., Sun, H., Liu, Z. H., Andrukhov, O., & Rausch-Fan, X. (2012). Effect of simvastatin on the osteogenic behavior of alveolar osteoblasts and periodontal ligament cells. *Human Cell*, 25, 29–35. <https://doi.org/10.1007/s13577-011-0028-x>
- Loos, B. G., Papantonopoulos, G., Jepsen, S., & Laine, M. L. (2015). What is the Contribution of Genetics to Periodontal Risk? *Dental Clinics of North America*, 59, 761–780. <https://doi.org/10.1016/j.cden.2015.06.005>
- Lopez, N. J., Quintero, A., Casanova, P. A., Ibieta, C. I., Baelum, V., & Lopez, R. (2012). Effects of periodontal therapy on systemic markers of inflammation in patients with metabolic syndrome: A controlled clinical trial. *Journal of Periodontology*, 83, 267–278. <https://doi.org/10.1902/jop.2011.110227>
- Lu, B., Parker, D., & Eaton, C. B. (2008). Relationship of periodontal attachment loss to peripheral vascular disease: An analysis of NHANES 1999–2002 data. *Atherosclerosis*, 200, 199–205. <https://doi.org/10.1016/j.atherosclerosis.2007.12.037>
- Luan, Z., Chase, A. J., & Newby, A. C. (2003). Statins inhibit secretion of metalloproteinases-1, -2, -3, and -9 from vascular smooth muscle cells and macrophages. *Arteriosclerosis, Thrombosis, and Vascular Biology*, 23, 769–775. <https://doi.org/10.1161/01.ATV.0000068646.76823.AE>
- Madrid, C., & Sanz, M. (2009). What influence do anticoagulants have on oral implant therapy? A systematic review. *Clinical Oral Implants Research*, 20(Suppl 4), 96–106. <https://doi.org/10.1111/j.1600-0501.2009.01770.x>
- Mahendra, J., Mahendra, L., Felix, J., & Romanos, G. (2013). Prevalence of periodontopathogenic bacteria in subgingival biofilm and atherosclerotic plaques of patients undergoing coronary revascularization surgery. *Journal of Indian Society of Periodontology*, 17, 719724. <https://doi.org/10.4103/0972-124X.124476>
- Matthews, J. B., Wright, H. J., Roberts, A., Cooper, P. R., & Chapple, I. L. (2007a). Hyperactivity and reactivity of peripheral blood neutrophils in chronic periodontitis. *Clinical and Experimental Immunology*, 147, 255–264. <https://doi.org/10.1111/j.1365-2249.2006.03276.x>
- Matthews, J. B., Wright, H. J., Roberts, A., Ling-Mountford, N., Cooper, P. R., & Chapple, I. L. (2007b). Neutrophil hyper-responsiveness in periodontitis. *Journal of Dental Research*, 86, 718–722. <https://doi.org/10.1177/154405910708600806>
- Meisel, P., Kroemer, H. K., Nauck, M., Holtfrete, B., & Kocher, T. (2014). Tooth loss, periodontitis, and statins in a population-based follow-up study. *Journal of Periodontology*, 85, e160–168. <https://doi.org/10.1902/jop.2013.130456>
- Mendez, M. V., Scott, T., LaMorte, W., Vokonas, P., Menzoian, J. O., & Garcia, R. (1998). An association between periodontal disease and peripheral vascular disease. *American Journal of Surgery*, 176, 153–157. [https://doi.org/10.1016/S0002-9610\(98\)00158-5](https://doi.org/10.1016/S0002-9610(98)00158-5)
- Miclotte, I., Vanhaverbeke, M., Agbaje, J. O., Legrand, P., Vanassche, T., Verhamme, P., & Politis, C. (2017). Pragmatic approach to manage new oral anticoagulants in patients undergoing dental extractions: A prospective case-control study. *Clinical Oral Investigations*, 21, 2183–2188. <https://doi.org/10.1007/s00784-016-2010-1>
- Minassian, C., D'Aiuto, F., Hingorani, A. D., & Smeeth, L. (2010). Invasive dental treatment and risk for vascular events: A self-controlled case series. *Annals of Internal Medicine*, 153, 499–506. <https://doi.org/10.7326/0003-4819-153-8-201010190-00006>
- Montenegro, M. M., Ribeiro, I. W. J., Kampits, C., Saffi, M. A. L., Furtado, M. V., Polanczyk, C. A., ... Rosing, C. K. (2019). Randomized controlled trial of the effect of periodontal treatment on cardiovascular risk biomarkers in patients with stable coronary artery disease: Preliminary findings of 3 months. *Journal of Clinical Periodontology*, 46, 321–331. <https://doi.org/10.1111/jcpe.13085>
- Mundy, G., Garrett, R., Harris, S., Chan, J., Chen, D., Rossini, G., ... Gutierrez, G. (1999). Stimulation of bone formation in vitro and in rodents by statins. *Science*, 286, 1946–1949. <https://doi.org/10.1126/science.286.5446.1946>
- Muniz, F., Taminski, K., Cavagni, J., Celeste, R. K., Weidlich, P., & Rosing, C. K. (2018). The effect of statins on periodontal treatment—a systematic review with meta-analyses and meta-regression. *Clinical Oral Investigations*, 22, 671–687. <https://doi.org/10.1007/s00784-018-2354-9>
- Munz, M., Richter, G. M., Loos, B. G., Jepsen, S., Divaris, K., Offenbacher, S., ... Schaefer, A. S. (2018). Genome-wide association meta-analysis of coronary artery disease and periodontitis reveals a novel shared risk locus. *Scientific Reports*, 8, 13678. <https://doi.org/10.1038/s41598-018-31980-8>
- Napenas, J. J., Hong, C. H., Brennan, M. T., Furney, S. L., Fox, P. C., & Lockhart, P. B. (2009). The frequency of bleeding complications after invasive dental treatment in patients receiving single and dual antiplatelet therapy. *Journal of the American Dental Association*, 140, 690–695. <https://doi.org/10.14219/jada.archive.2009.0255>
- Nathwani, S., & Martin, K. (2016). Exodontia in dual antiplatelet therapy: The evidence. *British Dental Journal*, 220, 235–238. <https://doi.org/10.1038/sj.bdj.2016.173>
- Nordendahl, E., Kjellstrom, B., Fored, C. M., Ekblom, A., Svensson, T., Norhammar, A., & Gustafsson, A. (2018). Invasive dental treatment and risk for a first myocardial infarction. *Journal of Dental Research*, 97, 1100–1105. <https://doi.org/10.1177/0022034518767834>
- O'Donnell, M. J., Chin, S. L., Rangarajan, S., Xavier, D., Liu, L., Zhang, H., ... INTERSTROKE Investigators (2016). Global and regional effects of potentially modifiable risk factors associated with acute stroke in 32 countries (INTERSTROKE): A case-control study. *Lancet*, 388, 761–775. [https://doi.org/10.1016/S0140-6736\(16\)30506-2](https://doi.org/10.1016/S0140-6736(16)30506-2)
- Offenbacher, S., Beck, J. D., Moss, K., Mendoza, L., Paquette, D. W., Barrow, D. A., ... Genco, R. J. (2009). Results from the Periodontitis and Vascular Events (PAVE) Study: A pilot multicentered, randomized, controlled trial to study effects of periodontal therapy in a secondary prevention model of cardiovascular disease. *Journal of Periodontology*, 80, 190–201. <https://doi.org/10.1902/jop.2009.080007>
- Orlandi, M., Graziani, F., & D'Aiuto, F. (2019). Periodontal therapy and cardiovascular risk. *Periodontology 2000*, (in press).
- Paraskevas, S., Huizinga, J. D., & Loos, B. G. (2008). A systematic review and meta-analyses on C-reactive protein in relation to periodontitis. *Journal of Clinical Periodontology*, 35, 277–290. <https://doi.org/10.1111/j.1600-051X.2007.01173.x>
- Park, S. Y., Kim, S. H., Kang, S. H., Yoon, C. H., Lee, H. J., Yun, P. Y., ... Chae, I. H. (2019). Improved oral hygiene care attenuates the cardiovascular risk of oral health disease: A population-based study from Korea.

- European Heart Journal*, 40, 1138–1145. <https://doi.org/10.1093/eurheartj/ehy836>
- Patel, J. P., Woolcombe, S. A., Patel, R. K., Obisesan, O., Roberts, L. N., Bryant, C., & Arya, R. (2017). Managing direct oral anticoagulants in patients undergoing dentoalveolar surgery. *British Dental Journal*, 222, 245–249. <https://doi.org/10.1038/sj.bdj.2017.165>
- Patil, V. A., & Desai, M. H. (2013). Effect of periodontal therapy on serum C-reactive protein levels in patients with gingivitis and chronic periodontitis: A clinicobiochemical study. *Journal of Contemporary Dental Practice*, 14, 233–237. <https://doi.org/10.5005/jp-journals-10024-1305>
- Paumelle, R., Blanquart, C., Briand, O., Barbier, O., Duhem, C., Woerly, G., ... Staels, B. (2006). Acute antiinflammatory properties of statins involve peroxisome proliferator-activated receptor- α via inhibition of the protein kinase C signaling pathway. *Circulation Research*, 98, 361–369. <https://doi.org/10.1161/01.RES.0000202706.70992.95>
- Perry, D. J., Noakes, T. J., Helliwell, P. S., & British Dental, S. (2007). Guidelines for the management of patients on oral anticoagulants requiring dental surgery. *British Dental Journal*, 203, 389–393. <https://doi.org/10.1038/bdj.2007.892>
- Poston, C. J., Pierce, T. C., Li, Y., Brinson, C. W., Lu, Z., Lauer, A. W., ... Huang, Y. (2016). Statin intake is associated with MMP-1 level in gingival crevicular fluid of patients with periodontitis. *Oral Diseases*, 22, 438–444. <https://doi.org/10.1111/odi.12474>
- Quist-Paulsen, P. (2010). Statins and inflammation: An update. *Current Opinion in Cardiology*, 25, 399–405. <https://doi.org/10.1097/HCO.0b013e3283398e53>
- Rafferty, B., Jönsson, D., Kalachikov, S., Demmer, R. T., Nowygrod, R., Elkind, M. S., ... Kozarov, E. (2011). Impact of monocytic cells on recovery of uncultivable bacteria from atherosclerotic lesions. *Journal of Internal Medicine*, 270, 273–280. <https://doi.org/10.1111/j.1365-2796.2011.02373.x>
- Range, H., Labreuche, J., Louedec, L., Rondeau, P., Planes, C., Sebbag, U., ... Meilhac, O. (2014). Periodontal bacteria in human carotid atherothrombosis as a potential trigger for neutrophil activation. *Atherosclerosis*, 236, 448–455. <https://doi.org/10.1016/j.atherosclerosis.2014.07.034>
- Reichert, S., Schulz, S., Benten, A. C., Lutze, A., Seifert, T., Schlitt, M., ... Schlitt, A. (2016). Periodontal conditions and incidence of new cardiovascular events among patients with coronary vascular disease. *Journal of Clinical Periodontology*, 43, 918–925. <https://doi.org/10.1111/jcpe.12611>
- Reyes, L., Herrera, D., Kozarov, E., Roldan, S., & Progulske-Fox, A. (2013). Periodontal bacterial invasion and infection: Contribution to atherosclerotic pathology. *Journal of Clinical Periodontology*, 40, S30–S50. <https://doi.org/10.1111/jcpe.12079>
- Rosenson, R. S., Tangney, C. C., & Casey, L. C. (1999). Inhibition of proinflammatory cytokine production by pravastatin. *Lancet*, 353, 983–984. [https://doi.org/10.1016/S0140-6736\(98\)05917-0](https://doi.org/10.1016/S0140-6736(98)05917-0)
- Roth, G. A., Forouzanfar, M. H., Moran, A. E., Barber, R., Nguyen, G., Feigin, V. L., ... Murray, C. J. (2015). Demographic and epidemiologic drivers of global cardiovascular mortality. *New England Journal of Medicine*, 372, 1333–1341. <https://doi.org/10.1056/NEJMoa1406656>
- Roth, G. A., Johnson, C., Abajobir, A., Abd-Allah, F., Abera, S. F., Abyu, G., ... Murray, C. (2017). Global, regional, and national burden of cardiovascular diseases for 10 causes, 1990 to 2015. *Journal of the American College of Cardiology*, 70, 1–25. <https://doi.org/10.1016/j.jacc.2017.04.052>
- Saffi, M. A. L., Rabelo-Silva, E. R., Polanczyk, C. A., Furtado, M. V., Montenegro, M. M., Ribeiro, I. W. J., ... Haas, A. N. (2018). Periodontal therapy and endothelial function in coronary artery disease: A randomized controlled trial. *Oral Diseases*, 24, 1349–1357. <https://doi.org/10.1111/odi.12909>
- Sakoda, K., Yamamoto, M., Negishi, Y., Liao, J. K., Node, K., & Izumi, Y. (2006). Simvastatin decreases IL-6 and IL-8 production in epithelial cells. *Journal of Dental Research*, 85, 520–523. <https://doi.org/10.1177/154405910608500608>
- Sangwan, A., Tewari, S., Singh, H., Sharma, R. K., & Narula, S. C. (2013). Periodontal status and hyperlipidemia: Statin users versus non-users. *Journal of Periodontology*, 84, 3–12. <https://doi.org/10.1902/jop.2012.110756>
- Sangwan, A., Tewari, S., Singh, H., Sharma, R. K., & Narula, S. C. (2016). Effect of hyperlipidemia on response to nonsurgical periodontal therapy: Statin users versus nonusers. *European Journal of Dentistry*, 10, 69–76. <https://doi.org/10.4103/1305-7456.175685>
- Saver, B. G., Hujoel, P. P., Cunha-Cruz, J., & Maupome, G. (2007). Are statins associated with decreased tooth loss in chronic periodontitis? *Journal of Clinical Periodontology*, 34, 214–219. <https://doi.org/10.1111/j.1600-051X.2006.01046.x>
- Saxlin, T., Suominen-Taipale, L., Knuuttila, M., Alha, P., & Ylostalo, P. (2009). Dual effect of statin medication on the periodontium. *Journal of Clinical Periodontology*, 36, 997–1003. <https://doi.org/10.1111/j.1600-051X.2009.01484.x>
- Schaefer, A. S., Bochenek, G., Jochens, A., Ellinghaus, D., Dommisch, H., Guzeldemir-Akcanat, E., ... Schreiber, S. (2015). Genetic evidence for PLASMINOGEN as a shared genetic risk factor of coronary artery disease and periodontitis. *Circulation: Cardiovascular Genetics*, 8, 159–167. <https://doi.org/10.1161/CIRCGENETICS.114.000554>
- Schaefer, A. S., Richter, G. M., Dommisch, H., Reinartz, M., Nothnagel, M., Noack, B., ... Schreiber, S. (2011). CDKN2BAS is associated with periodontitis in different European populations and is activated by bacterial infection. *Journal of Medical Genetics*, 48, 38–47. <https://doi.org/10.1136/jmg.2010.078998>
- Schenkein, H. A., & Loos, B. G. (2013). Inflammatory mechanisms linking periodontal diseases to cardiovascular diseases. *Journal of Clinical Periodontology*, 40(Suppl 14), S51–69. <https://doi.org/10.1111/jcpe.12060>
- Schenkein, H. A., Papapanou, P. N., Genco, R., & Sanz, M. (2019). Mechanisms underlying the association between periodontitis and atherosclerotic disease. *Periodontology 2000*, (in press).
- Sen, S., Giamberardino, L. D., Moss, K., Morelli, T., Rosamond, W. D., Gottesman, R. F., ... Offenbacher, S. (2018). Periodontal disease, regular dental care use, and incident ischemic stroke. *Stroke*, 49, 355–362. <https://doi.org/10.1161/STROKEAHA.117.018990>
- Sen, S., Sumner, R., Hardin, J., Barros, S., Moss, K., Beck, J., & Offenbacher, S. (2013). Periodontal disease and recurrent vascular events in stroke/transient ischemic attack patients. *Journal of Stroke and Cerebrovascular Diseases*, 22, 1420–1427. <https://doi.org/10.1016/j.jstrokecerebrovasdis.2013.06.024>
- Sharma, P., Dietrich, T., Ferro, C. J., Cockwell, P., & Chapple, I. L. (2016). Association between periodontitis and mortality in stages 3–5 chronic kidney disease: NHANES III and linked mortality study. *Journal of Clinical Periodontology*, 43, 104–113. <https://doi.org/10.1111/jcpe.12502>
- Shi, Q., Xu, J., Zhang, T., Zhang, B., & Liu, H. (2017). Post-operative bleeding risk in dental surgery for patients on oral anticoagulant therapy: A meta-analysis of observational studies. *Frontiers in Pharmacology*, 8, 58. <https://doi.org/10.3389/fphar.2017.00058>
- Sixth Joint Task Force of the European Society of Cardiology and Other Societies on Cardiovascular Disease Prevention in Clinical Practice (2016) European guidelines on CVD prevention in clinical practice 2016. *European Journal of Preventive Cardiology*, 23, NP1–NP96. <https://doi.org/10.1177/2047487316653709>
- Smeeth, L., Thomas, S. L., Hall, A. J., Hubbard, R., Farrington, P., & Vallance, P. (2004). Risk of myocardial infarction and stroke after acute infection or vaccination. *New England Journal of Medicine*, 351, 2611–2618. <https://doi.org/10.1056/NEJMoa041747>

- Steffel, J., Verhamme, P., Potpara, T. S., Albaladejo, P., Antz, M., Desteghe, L.... ESC Scientific Document Group (2018a). The 2018 European Heart Rhythm Association Practical Guide on the use of non-vitamin K antagonist oral anticoagulants in patients with atrial fibrillation: Executive summary. *Europace*, 20, 1231–1242. <https://doi.org/10.1093/europace/euy054>
- Steffel, J., Verhamme, P., Potpara, T. S., Albaladejo, P., Antz, M., Desteghe, L.... ESC Scientific Document Group (2018b). The 2018 European Heart Rhythm Association Practical Guide on the use of non-vitamin K antagonist oral anticoagulants in patients with atrial fibrillation. *European Heart Journal*, 39, 1330–1393. <https://doi.org/10.1093/eurheartj/ehy136>
- Subramanian, S., Emami, H., Vucic, E., Singh, P., Vijayakumar, J., Fifer, K. M., ... Tawakol, A. (2013). High-dose atorvastatin reduces periodontal inflammation: A novel pleiotropic effect of statins. *Journal of the American College of Cardiology*, 62, 2382–2391. <https://doi.org/10.1016/j.jacc.2013.08.1627>
- Teeuw, W. J., Laine, M. L., Bizzarro, S., & Loos, B. G. (2015). A lead ANRIL polymorphism is associated with elevated CRP levels in periodontitis: A pilot case-control study. *PLoS ONE*, 10, e0137335. <https://doi.org/10.1371/journal.pone.0137335>
- Teeuw, W. J., Slot, D. E., Susanto, H., Gerdes, V. E., Abbas, F., D'Aiuto, F., ... Loos, B. G. (2014). Treatment of periodontitis improves the atherosclerotic profile: A systematic review and meta-analysis. *Journal of Clinical Periodontology*, 41, 70–79. <https://doi.org/10.1111/jcpe.12171>
- Ting, M., Whitaker, E. J., & Albandar, J. M. (2016). Systematic review of the in vitro effects of statins on oral and perioral microorganisms. *European Journal of Oral Sciences*, 124, 4–10. <https://doi.org/10.1111/eos.12239>
- Tomas, I., Diz, P., Tobias, A., Scully, C., & Donos, N. (2012). Periodontal health status and bacteraemia from daily oral activities: Systematic review/meta-analysis. *Journal of Clinical Periodontology*, 39, 213–228. <https://doi.org/10.1111/j.1600-051X.2011.01784.x>
- Tonetti, M. S., Van Dyke, T. E. & Working group 1 of the Joint EFP/AAP Workshop (2013). Periodontitis and atherosclerotic cardiovascular disease: Consensus report of the Joint EFP/AAP Workshop on Periodontitis and Systemic Diseases. *Journal of Clinical Periodontology*, 40(Suppl 14), S24–29. <https://doi.org/10.1111/jcpe.12089>
- U.S. Senate Committee on Health, Education Labor & Pensions (2011). *The state of chronic disease prevention: Hearing before the Committee on Health, Education, Labor, and Pensions*.
- Valgimigli, M., Bueno, H., Byrne, R. A., Collet, J. P., Costa, F., Jeppsson, A.... ESC Scientific Document Group (2018). 2017 ESC focused update on dual antiplatelet therapy in coronary artery disease developed in collaboration with EACTS. *European Journal of Cardio-Thoracic Surgery*, 53, 34–78. <https://doi.org/10.1093/ejcts/ezx334>
- Velsko, I. M., Chukkapalli, S. S., Rivera, M. F., Lee, J. Y., Chen, H., Zheng, D., ... Kesavalu, L. (2014). Active invasion of oral and aortic tissues by *Porphyromonas gingivalis* in mice causally links periodontitis and atherosclerosis. *PLoS ONE*, 9, e97811. <https://doi.org/10.1371/journal.pone.0097811>
- Velsko, I. M., Chukkapalli, S. S., Rivera-Kweh, M. F., Zheng, D., Aukhil, I., Lucas, A. R., ... Kesavalu, L. (2015). Periodontal pathogens invade gingiva and aortic adventitia and elicit inflammasome activation in v6 integrin-deficient mice. *Infection and Immunity*, 83, 4582–4593. <https://doi.org/10.1128/IAI.01077-15>
- Vidal, F., Cordovil, I., Figueredo, C. M., & Fischer, R. G. (2013). Non-surgical periodontal treatment reduces cardiovascular risk in refractory hypertensive patients: A pilot study. *Journal of Clinical Periodontology*, 40, 681–687. <https://doi.org/10.1111/jcpe.12110>
- Viereck, V., Grundker, C., Blaschke, S., Frosch, K. H., Schoppet, M., Emons, G., & Hofbauer, L. C. (2005). Atorvastatin stimulates the production of osteoprotegerin by human osteoblasts. *Journal of Cellular Biochemistry*, 96, 1244–1253. <https://doi.org/10.1002/jcb.20598>
- Wilkins, E., Wilson, L., Wickramasinghe, K., Bhatnagar, P., Leal, J., Luengo-Fernandez, R., ... Townsend, N. (2017). *European Cardiovascular Disease Statistics 2017*. Brussels: European Heart Network.
- Wright, H. J., Matthews, J. B., Chapple, I. L., Ling-Mountford, N., & Cooper, P. R. (2008). Periodontitis associates with a type 1 IFN signature in peripheral blood neutrophils. *Journal of Immunology*, 181, 5775–5784. <https://doi.org/10.4049/jimmunol.181.8.5775>
- Yagyu, T., Kawakami, M., Ueyama, Y., Imada, M., Kurihara, M., Matsusue, Y., ... Kirita, T. (2017). Risks of postextraction bleeding after receiving direct oral anticoagulants or warfarin: A retrospective cohort study. *British Medical Journal Open*, 7, e015952. <https://doi.org/10.1136/bmjopen-2017-015952>
- Yang, J., Wu, J., Liu, Y., Huang, J., Lu, Z., Xie, L., ... Ji, Y. (2014). *Porphyromonas gingivalis* infection reduces regulatory T cells in infected atherosclerosis patients. *PLoS ONE*, 9, e86599. <https://doi.org/10.1371/journal.pone.0086599>
- Yang, S., Shi, Q., Liu, J., Li, J., & Xu, J. (2016). Should oral anticoagulant therapy be continued during dental extraction? A meta-analysis. *BMC Oral Health*, 16, 81. <https://doi.org/10.1186/s12903-016-0278-9>
- Yang, S., Zhao, L., Cai, C., Shi, Q., Wen, N., & Xu, J. (2018). Association between periodontitis and peripheral artery disease: A systematic review and meta-analysis. *BMC Cardiovascular Disorders*, 18, 141. <https://doi.org/10.1186/s12872-018-0879-0>
- Yebo, H. G., Aschmann, H. E., Kaufmann, M., & Puhan, M. A. (2019). Comparative effectiveness and safety of statins as a class and of specific statins for primary prevention of cardiovascular disease: A systematic review, meta-analysis, and network meta-analysis of randomized trials with 94,283 participants. *American Heart Journal*, 210, 18–28. <https://doi.org/10.1016/j.ahj.2018.12.007>
- Zhou, Q. B., Xia, W. H., Ren, J., Yu, B. B., Tong, X. Z., Chen, Y. B., ... Yang, J. Y. (2017). Effect of intensive periodontal therapy on blood pressure and endothelial microparticles in patients with prehypertension and periodontitis: A randomized controlled trial. *Journal of Periodontology*, 88, 711–722. <https://doi.org/10.1902/jop.2017.160447>

SUPPORTING INFORMATION

Additional supporting information may be found online in the Supporting Information section at the end of the article.

How to cite this article: Sanz M, Marco del Castillo A, Jepsen S, et al. Periodontitis and cardiovascular diseases: Consensus report. *J Clin Periodontol*. 2019;00:1–17. <https://doi.org/10.1111/jcpe.13189>